

8 Literature Cited and General References

- Aber, J.D. 1992. Nitrogen cycling and nitrogen saturation in temperate forest ecosystems. In: TREE. 7(7):220-224.
- Aber, J.D. 1993. Modification of nitrogen cycling at the regional scale: the subtle effects of atmospheric deposition. pp. 163-174. In: McDonnell, M.J. and S.T.A. Pickett (eds.), Humans as components of ecosystems: the ecology of subtle human effects and populated areas. Springer-Verlag, N.Y. 364 pp.
- Aber, J.D., D.B. Botkin, and J.M. Melillo. 1978. Predicting the effects of different harvesting regimes on forest floor dynamics in northern hardwoods. Can. J. For. Res. Vol 8: 306-315
- Aber, J.D., J. Knute, P.S. Nadelhoffer, P. Steudler, and J.M. Melillo. 1989. Nitrogen saturation in northern forest ecosystems: excess nitrogen from fossil fuel combustion may stress the biosphere. Bioscience. 39(6):378-386.
- Aber, J.D., J.M. Melillo, and C.A. McClaugherty. 1989. Predicting long-term patterns of mass loss, nitrogen dynamics, and soil organic matter formation from initial fine litter chemistry in temperate forest ecosystems. Canadian Journal of Botany. 68:2201-2208.
- Aber, J.D., J.M. Melillo, K.J. Nadelhoffer, J. Pastor, and R.D. Boone. 1991. Factors controlling nitrogen cycling and nitrogen saturation in northern temperate forest ecosystems. Ecological Applications, 1(3), pp. 303-315.
- Agren, G.I. and E. Bosatta. 1988. Nitrogen saturation of terrestrial ecosystems. Environmental Pollution, 54: 185-197.
- Aiken, G.R., Capel, P.D., Furlong, E.T., Hult, M.F., and Thorn, K.A., 1991. Mechanisms controlling the transport of organic chemicals in subsurface environments, in Mallard, G.E., and Aronson, D.A., eds., U.S. Geological Survey Toxic Substances Hydrology Program--Proceedings of the technical meeting, Monterey, California, March 11-15, 1991: U.S. Geological Survey Water-Resources Investigations Report 91-4034, p. 633-637
- Alverson, W.S., and D.M. Waller. 1997. Deer populations and the widespread failure of hemlock regeneration in northern forests. In: The Science of Overabundance: deer ecology and population management. Smithsonian Institution Press. 402pp.
- American Forest Council. 1986. Your water - a forest product. GreenAmerica, American Forest Council, 1250 Connecticut Ave., NW, Washington, D.C. 20036
- Anacostia Restoration Team. 1992. Watershed restoration sourcebook. In: Collected papers presented at the conference: Restoring our home river: water quality and habitat in the Anacostia. Nov. 6 and 7. College Park, MD.
- Angelstam, P., P. Wikberg, P. Danilov, W.E. Faber, and K. Nygren. 2000. Effects of moose density on timber quality and biodiversity restoration in Sweden, Finland, and Russian Karelia. Alces. 36: 133-145.
- Apple, L.L. 1983. The use of beavers in riparian/aquatic habitat restoration in a cold desert, gully-cut stream system: a case history. Proceeding of the 18th Annual Meeting American Fisheries Society, Laramie, Wyoming, pages 29-35.
- Apple, L.L. no date. Riparian habitat restoration in cold desert, gully-cut stream systems: an innovative, cost effective, ecological approach. unpubl. manuscript. U.S. Bureau of Land Management, Rock Springs, Wyoming.

- Arbogast, C., Jr. 1985. Marking guides for northern hardwoods under the selection system. USDA Forest Service, Lake States Forest Experiment Station Paper 56, 19 pp.
- Armstrong, G.W. 1999. A stochastic characterization of a natural disturbance regime of the boreal mixed wood forest with implications for sustainable forest management. *Can. J. For. Res* 29(4): 424-433.
- Arora, K. S.K. Michelson, J.L. Baker, D.P. Tierney, and C.J. Peters. 1996. Herbicide retention by vegetative buffer strips from runoff under natural rainfall. *Transactions of the American Society of Agricultural Engineers* 39(6):2155-2162.
- Art, H.W. and D.P. Dethier. 1986. Influence of vegetative succession on soil chemistry of the Berkshires. Water Resources Research Center Publication, University of Massachusetts, Amherst, Project No. G912-05.
- Attiwill, P.M. 1994a. The disturbance of forest ecosystems: the ecological basis for conservative management. *Forest Ecology and Management*, 63:247-300.
- Attiwill, P.M. 1994b. Ecological disturbance and the conservative management of eucalypt forests in Australia. *Forest Ecology and Management*, 63:301-346.
- Austin, P., R. Bishop, D. Chandler, P. Deslauriers, and M. Hopkinson. 1992. 1992 water quality report: Quabbin Reservoir and Ware River watersheds. Metropolitan District Commission, Division of Watershed Management. 120 pp.
- Austin, P., R. Bishop, D. Chandler, and P. Deslauriers. 1990. 1990 water quality report: Quabbin Reservoir and Ware River watersheds. Metropolitan District Commission, Division of Watershed Management. 89 pp.
- Baker, John H. 1984. Sulfate retention by Forest Soils of Central New England. Publication No. 151, Water Resources Research Center, University of Massachusetts at Amherst.
- Banks, C.C. 1973. The strength of trees. *Journal of the Institute for Wood Science*, 6:44-50.
- Barnes, D. M., and A. U. Malik. 2001. Effects of beaver (*Castor canadensis*) herbivory on streamside vegetation in a northern Ontario watershed. *Canad. Field Nat.* 115.
- Barnes, William J., and Eric Dribble. 1988. The effects of beaver on riverbank forest succession. *Can. J. Bot.* 66:40-44.
- Barten, P.K. 2006. "Water in the Woods", notes from a practical seminar delivered to professional foresters.
- Batchelder, G.L., Yuretich, R.F. and W.C. Leonard. 1983. Hydrogeochemical cycling in a watershed-groundwater system: Fort River and Cadwell Creek, Massachusetts. Water Resources Research Center, University of Massachusetts. Pub. No. 145. 80 pp.
- Benton, C., F. Khan, P. Monaghan, W.N. Richards and C.B. Shedden. 1983. The contamination of a major water supply by gulls (*Larus* sp.): a study of the problem and remedial action taken. *Water Res.* 17:789-798.
- birds in southern New England. *J. Wildl. Manage.* 50(3):398-405.
- Beschta, R.L. 1990. Effects of fire on water quantity and quality. In: Walstad, J.D., S.R. Radosevich, D.V. Sandberg, eds. *Natural and prescribed fire in Pacific Northwest forests*. Corvallis, OR. Oregon State University Press. 219-232.
- Bishop, R.P. 1992. Quabbin Reservation sanitary district: a survey of potential water pollution sources. Metropolitan District Commission, Division of Watershed Management.
- Blackburn, P. and J.A. Petty. 1988. Theoretical calculations of the influence of spacing on stand stability. *Forestry*, 61: 235-244

- Blodgett, B. G. and R.A. Clark. no date. A check list for birds of the Quabbin Reservation, west central Massachusetts. Friends of Quabbin, Belchertown.
- Blodgett, D. 1985. Snag and den tree management guidelines. Habitat Highlights. Vol. 5. No. 4. 2pp.
- Bolgiano, C. 1989. A case for eastern old-growth. Amer. Forests. May-June:26-31,48.
- Bormann, F.H. and G.E. Likens. 1979a. Catastrophic disturbance and the steady state in Northern Hardwood Forests. American Scientist 67:660-669.
- Bormann, F.H. and G.E. Likens. 1979b. Patterns and process in a forested ecosystem; disturbance, development and the steady state based on the Hubbard Brook Ecosystem Study. Springer-Verlag. New York, N.Y. 253 pp.
- Bormann, F.H. and G.E. Likens. 1970. The nutrient cycles of an ecosystem: measurements taken in a New Hampshire forest. Scientific American. 223(4).
- Bormann, F.H., G.E. Likens, and J.S. Eaton. 1969. Biotic regulation of particulate and solution losses from a forest ecosystem. Bioscience Vol 19, July, pp 600-610.
- Bormann, F.H., G.E. Likens, D.W. Fisher, R.S. Pierce. 1968. Nutrient loss accelerated by clear-cutting of a forest. Ecosystem Science, February, pp 882-884.
- Bormann, F.H., G.E. Likens, T.G. Siccama, R.S. Pierce, and J.S. Eaton. 1974. The export of nutrients and recovery of stable conditions following deforestation at Hubbard Brook. Ecol. Monographs 44:255-277.
- Bowden, R.D., M.S. Castro, J.M. Melillo, P.A. Steudler and J.D. Aber. 1993. Fluxes of greenhouse gases between soils and the atmosphere in a temperate forest following a simulated hurricane blowdown. Biogeochemistry, 21:61-71.
- Bradshaw, F.J. 1992. Quantifying edge effects and patch size for multiple-use silviculture – a discussion paper. Forest Ecology and Management. 48:249-264.
- Brandner, T.A., R.O. Peterson, and K.L. Risenhoover. 1990. Balsam fir on Isle Royale: effects of moose herbivory and population density. Ecology. 71(1): 155-164.
- Branvold, D.K., C.J. Popp and J.A. Briery. 1976. Waterfowl refuge effect on water quality: II. chemical and physical parameters. Journal of Water Pollution Control Fed. 48(4):680-687.
- Brayton, D. Scott. 1984. The beaver and the stream. Journal of Soil Water Conservation. March/April: 108-109.
- Broadmeadow, M.S.J. (ed). 2001. Climate change and U.K. forests. Forestry Commission Bulletin #125. Forestry Commission, Edinburgh. 198 p.
- Bromley, P. T., J. Starr, J. Sims, and D. Coffman. 1997. A landowner's guide to wildlife abundance through forestry. University of Virginia Cooperative Extension. Publication: 420-138.
- Bromley, S.W. 1935. The original forest types of southern New England. Ecol. Monographs Vol 5(1), January, pp 63-89.
- Brooks, K.N., P.F. Ffolliott, H.M. Gregersen, and L.F. DeBano. 1997. Hydrology and the management of watersheds. 2nd Edition. Iowa State University Press. Ames, IA
- Brooks, R.T. and T.D. Kyker-Snowman. 2007. Forest floor temperature and relative humidity following timber harvesting in southern New England, USA. Forest Ecology and Management. Accepted July 21, 2007. *In press*.
- Brown, G.W. 1976. Forestry and water quality. School of Forestry, Oregon State University, Corvallis, Oregon. Reprinted by the Oregon State University Book Stores, Inc.

- Brown, K.A., P.H. Freer-Smith, G.D. Howells, R.A. Skeffington, and R.B. Wilson. 1988. Rapporteurs report on discussions at the workshop on excess nitrogen deposition, Leatherhead, September 1987. *Environmental Pollution*, 54:285-295.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. *Distance sampling: estimating abundance of biological populations*. Chapman & Hall, London. 446 pp.
- Burns, R.G. and J.D. Hewlett. 1983. A decision model to predict sediment yield from forest practices. *Water Resources Bulletin* Vol 19, No 1, pp 9-14.
- Burns, R.M. and B.H. Honkala, tech. coords. 1990. *Silvics of North America: Vol.1. Conifers, Vol.2. Hardwoods*. Agricultural Handbook No.654. U.S. Department of Agriculture, Forest Service, Washington, D.C. Vol.1. 675 pp.; Vol.2. 877 pp.
- Buzzell, G. 1991. 1990 Wachusett storm salvage study. MDC Publication.
- Buzzell, G. 1993. Study of regeneration response at Walnut Hill, Sudbury Reservoir watershed. Unpublished MDC report.
- Carlson, J. 1994. Effects of a simulated hurricane on downed woody fuel loading in a central Massachusetts hardwood stand. Abstr. 5th Ann. Harvard Forest Ecology Symp., April 25.
- Carlton, M. 1990. Literature review: water quality implications of converting forested watersheds to principally herbaceous cover - implications to the Quabbin watershed. MDC Publication.
- Carroll, C.F. 1973. *The timber economy of puritan New England*. Brown University Press, Providence, RI. Pp.221
- Chadwick, N.L., D. R. Progulske, and J. T. Finn. 1986. Effects of fuelwood cutting on
- Chang, M. 2003. *Forest hydrology: an introduction to water and forests*. CRC Press LLC, Boca Raton, FL. 373 p.
- Chen, H. and T.J. Cutright. 2003. Preliminary evaluation of microbially mediated precipitation of cadmium, chromium, and nickel by rhizosphere consortium. *Journal of Environmental Engineering – ASCE*. 129(1):4-9.
- Christ, M.J., W.T. Peterjohn, J.R. Cumming, and M.B. Adams. 2002. Nitrification potentials and landscape, soil and vegetation characteristics in two central Appalachian watersheds differing in NO₃- export. *For Eco Manage* 159: 145-158.
- Clark, D. E. 1999. Quabbin and Wachusett Reservoir Watersheds Aquatic Wildlife Pathogen Control Zones. Metropolitan District Commission, Division of Watershed Management, Natural Resources Section. 17pp.
- Clark, D.E. 2004. Quabbin Reservation White-Tailed Deer Impact Management Program: 5-Year (2000-2004) Review and New 5-Year Plan (2005-2009). Dept. Cons. And Recreation. Div. of Water Supply Prot.
- Cline, A.C. 1939. The restoration of watershed forests in the hurricane area. *Journal of the New England Water Works Association*. 53(2):223-237.
- Cline, A.C. and S.H. Spurr. 1942. The virgin upland forest of central New England: a study of old growth stands in the Pisgah Mountain section of southwestern New Hampshire. *Harvard Forest Bulletin* No 21, 58 pp.
- Connecticut Department of Environmental Protection. 1991. Guidelines for enhancing Connecticut's wildlife habitat through forestry operations. Dept. Environ. Prot. Wildl. Div. 14pp.
- Connor, K.J., W.B. Ballard, T. Dilworth, S. Mahoney, and D. Anions. 2000. Changes in structure of a boreal forest community following intense herbivory by moose. *Alces*. 36: 111-132.

- Cooper-Ellis, S., D.R. Foster, G. Carlton, and A. Lezberg. 1999. Forest response to catastrophic wind: results from an experimental hurricane. *Ecology* 80(8): 2683-2696.
- Corbett, E.S. and W. Spencer. 1975. Effects of management practices on water quality and quantity: Baltimore, Maryland municipal watersheds. In: *Municipal Watershed Management Symposium Proceedings*, USDA Forest Service General Technical Report NE-13, pgs. 25-31.
- Cosby, B.J., Wright, R.F., Hornberger, G.M., and J.N. Galloway. 1985. Modeling the effects of acid deposition: estimation of long-term water quality responses in a small forested catchment. *Water Resources Research* 21(11):1591-1601.
- Crabtree, R.C. and F.A. Bazzaz. 1993. Seedling response of four birch species to simulated nitrogen deposition: ammonium vs. nitrate. *Ecological Applications* 3(2) pp. 315-321.
- Craun, G.F. 1984. Waterborne outbreaks of giardiasis, current status. IN: Erlandsen and Meyer (eds.). *Giardia and Giardiasis*. Plenum Press. N.Y. pgs.. 243-261.
- Cronin, W. 1983. *Changes in the land: Indians, colonists, and the ecology of New England*. Hill and Wang. New York, NY. 241 pp.
- Curran, M.L. and D. F. Dincauze. 1977. Paleo-Indians and paleo-lakes: new data from the Connecticut drainage. *Annals of the New York Academy of Sciences*. 288:333-348.
- Currie, W. and J.D. Aber. 1994. Modeling litter decomposition and N dynamics across LTER sites as part of the LIDET study. Abstracts from the 5th Annual Harvard Forest Ecology Symposium. Petersham, MA.
- Currier, J.B., L.E. Siverts, and R.C. Maloney. 1979. An approach to water resources evaluation of non-point sources from silvicultural activities - a procedural handbook. In: *Best Management Practices for Agriculture and Silviculture: Proceedings of the 1978 Cornell Agricultural Waste Management Conference*, Ann Arbor Science Publishers Inc., Ann Arbor, Michigan
- Curtis, J.T. 1959. *The vegetation of Wisconsin*. Univ. Wisconsin Press, Madison. 657 pp.
- Danell, K., L.Edenius, and P.Lundberg. 1991. Herbivory and tree stand composition: moose patch use in winter. *Ecology*. 72(4): 1350-1357.
- Davies, R.B. and C.P. Hibler. 1976. Animal reservoirs and cross-species transmission of *Giardia*. pp. 104-126 IN: Jakubowski and Hoff (eds). *Waterborne transmission of giardiasis*. U.S. EPA - 600/9-79-001.
- Davis, M.B., 1969. Climatic changes in southern Connecticut recorded by pollen deposition at Rogers Lake. *Ecology* 50, 409-422.
- Davis, M.B., 1983. Holocene vegetational history of the eastern United States. In: Wright Jr., H.E. (Ed.), *Late Quaternary Environments of the United States: Volume 2, The Holocene*. University of Minnesota Press, Minneapolis, Minnesota, pp. 166-181.
- Davis, J.W., G.A. Goodwin, and R.A. Ockenfels. 1983. Snag habitat management: proceedings of the symposium. U.S. For. Serv. Gen. Tech. Rep. RM-99. 226pp.
- Day, G.M. 1953. The Indian as an ecological factor. *Ecology*. 34(2): 329-347.
- DeBano, L.F., D.G. Neary, and P. F. Ffolliott. 1998. *Fire's effects on ecosystems*. John Wiley Publishers, Inc., New York, NY. 333 p.
- DeCoster, L.A. and J. Herrington. 1988. Is a tree a heavy drinker, or does it just pump water? *The American Tree Farmer*, May-June, 1988.
- DeGraaf, R.M. and D.A. Richard. 1987. Forest wildlife of Massachusetts: cover type, size class and special habitat relationships. Univ. Mass. Coop. Ext. Serv. Publ. C-182.

- DeGraaf, R.M., and A.L. Shigo. 1985. Managing cavity trees for wildlife in the Northeast. U.S. For. Serv. Gen. Tech. Rep. NE-101. 21pp.
- DeGraaf, R.M., and D.D. Rudis. 1986. New England wildlife: habitat, natural history, and distribution. U.S. For. Serv. Gen. Tech. Rep. NE-108. 491pp.
- DeGraaf, R.M., and K.E. Evans. 1980. Management of north central and northeastern forests for nongame birds. U.S. For. Serv. Gen. Tech. Rep. NC-51. 275pp.
- DeGraaf, R.M., and W.M. Healy. 1988. Is forest fragmentation a management issue in the northeast? U.S. For. Serv. Gen. Tech. Rep. NE-140. 32pp.
- DeGraaf, R.M., M. Yamasaki, W.B. Leak, and J.W. Lanier. 1992. New England wildlife: management of forested habitats. U.S. For. Serv. Gen. Tech. Rep. NE-144. 271pp.
- De la Cretaz, A.L. and P.K. Barten. 2007. Land use effects on streamflow and water quality in the Northeastern United States. CRC Press. Boca Raton, FL. 319 pp.
- DeMaynadier, P. G., and M. L. Hunter, Jr. 2000. Road effects on amphibian movements in a forested landscape. *Nat. Areas. J.* 20:56-65.
- Dessecker, D.R., and D.G. McAuley. 2001. Importance of early successional habitat to ruffed grouse and American woodcock. *Wildl. Soc. Bull.* 29(2):456-465.
- Dettmers, R. and K. Rosenburg. 2000. Partners in flight landbird conservation plan: physiographic area 9: southern New England. Version 1.0. American Bird Conservancy. Website: http://www.blm.gov/wildlife/plan/pl_09_10.pdf
- Devito, K. J., and P. J. Dillion. 1993. Importance of runoff and winter anoxia to the P and N dynamics of a beaver pond. *Can. J. Fish. Aquat. Sci.* 50:2222-2244.
- Diesch, S.L. 1970. Disease transmission of water-borne organisms of animal origin. pp. 269-285 IN: T.L. Willrich and G.E. Smith (eds). *Agricultural practices and water quality.* Iowa St. Univ. Press.
- Dincauze, D.F. 1974. An introduction to archaeology in the greater Boston area. *Archaeology of Eastern North America.* 2(1):47.
- Dincauze, D.F. and M.T. Mulholland. 1977. Early and middle archaic site distributions and habitats in southern New England. *Annals of the New York Academy of Sciences.* 288:439-455.
- Dincauze, D.F.; P.A. Thomas; J.S. Wilson; and M.T. Mulholland. 1976. Cultural resource survey and impact evaluation report, state Route 2: Greenfield, Gill, Erving, Wendell, Orange, Franklin County, Massachusetts. University of Massachusetts, Amherst.
- Dissmeyer, G.E., ed. 2000. Drinking water from forests and grasslands: a synthesis of the scientific literature. Gen. Tech. Rep. SRS-39. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 246 p.
- Dobson, J.E., R.M. Rush, and R.W. Peplies. 1990. Forest blowdown and lake acidification. *Annals of the Association of American Geographers.* 80(3):343-361.
- Donkor, N.T., and J.M. Fryxell. 1999. Impact of beaver foraging on structure of lowland boreal forests of Algonquin Provincial Park, Ontario. *For. Ecol. Manage.* 118:83-92.
- Douglass, J.E. 1983. The potential for water yield augmentation from forest management in the eastern United States. *Water Resources Bulletin Vol 19(3),* pp 351-358.
- Douglas, J.E. and W.T. Swank. 1975. Effects of management practices on water quality and quantity: Coweeta Hydrologic Laboratory, North Carolina. In: *Municipal Watershed Management Symposium Proceedings, USDA Forest Service, General Technical Report, NE-13,* pp 1-12

- Douglass, J.E. and W.T. Swank. 1972. Streamflow modification through management of eastern forests. U.S.D.A. Forest Service, Research Paper SE-94, 15p.
- Driscoll, Charles T., G.E. Likens, L.O. Hedin, J. S. Eaton and F.H. Bormann. 1989. Changes in the chemistry of surface waters: Twenty-five year results at the Hubbard Brook Experimental Forest, NH. *Environ. Sci. Technol.* 23(2):137-143.
- Dunne, T. and L.B. Leopold. 1978. *Water in environmental planning*, W.H. Freeman and Company, New York, NY, 818 pp.
- Dyrness, C.T. 1965. Erodibility and erosion potential of forest watersheds. In: *Forest Hydrology: Proceedings of a National Science Foundation Advanced Science Seminar at Pennsylvania State University, University Park, PA, August 29-Sept 10, 1965*, Edited by W.E. Sopper and H.W. Lull, Pergamon Press, London, pp 559-611.
- Edenius, L. 1994. Foraging behavior by moose in forest stands subjected to severe browsing. *Alces*. 30: 37-40.
- Edwards, Nicole T., and David L. Otis. 1999. Avian communities and habitat relationships in South Carolina Piedmont beaver ponds. *Am. Midl. Nat.* 141:158-171.
- Elliot, C. A. 1988. A forester's guide to managing wildlife habitats in Maine. *U. Maine Coop. Ext.* 46pp.
- Erlandsen, S.L. and W. J. Bemrick. 1988. Waterborne giardiasis: sources of *Giardia* cysts and evidence pertaining to their implication in human infection. pp. 227-236 IN: P.M. Wallis and B.R. Hammond (eds). *Advances in Giardia research*. Univ. Calgary Press, Calgary, Alberta.
- Eschner, A.R. 1978. Long term observations and analyses of the hydrology of small, relatively natural watersheds in the Adirondack Mountains. Reprinted from: *Tagungsbericht Forschung in mitteleuropaischen Nationalparken. Schriftenreihe des Bayerischen Staatsministeriums fur Ernährung, Landwirtschaft und Forsten. Grafenau 25-27 Mai 1978.*
- Eschner, A.R. and D.L. Mader. 1975. Management implications of other hydrologic research in the northeast. In: *Municipal Watershed Management Symposium Proceedings*. USDA Forest Service: General Technical Report NE-13.
- Eschner, A.R. and D.R. Satterlund. 1965. Forest protection and streamflow from an Adirondack watershed. *Wat. Res. Res.* 2(4):765-783.
- Executive Office of Environmental Affairs, Commonwealth of Massachusetts. 2000. *The state of our environment*. 150 pp.
- February 5-7, 2002. Rutgers – the State University of New Jersey, East Brunswick, NJ. 403 p.
- Federal Register. 1987. 52(212):42178. Nov. 3.
- Federer, C.A. and D. Lash. 1978. Brook: a hydrologic simulation model for eastern forests. *Water Resource Research Center, University of New Hampshire, Durham, Research Report Number 19.*
- Fennell, H., D.B. James and J. Morris. 1974. Pollution of a storage reservoir by roosting gulls. *Water Treat. Exam.* 23:5-24.
- Finzi, A.C., C.D. Canham, and N. Van Breemen. 1998. Canopy tree-soil interactions within temperate forests: species effects on pH and cations. *Ecological Applications*, 8(2), pp. 447-454
- Flatebo, G., C. R. Foss, and S. K. Pelletier. 1999. Biodiversity in the forests of Maine: guidelines for land management. *Univ. Maine Coop. Ext. No.7147* 167pp.
- Forman, R.T.T. and M. Godron. 1986. *Landscape ecology*. Wiley & Sons, N.Y.
- Foster, D. R. 1999. *Thoreau's country*. Harvard University Press, Cambridge, MA.
- Foster, D.R. 1988a. Species and stand response to catastrophic wind in central New England, U.S.A. *Journal of Ecology* 76:135-151.

- Foster, D.R. 1988b. Disturbance history, community organization and vegetation dynamics of the old-growth Pisgah Forest, South-Western New Hampshire, U.S.A. *Journal of Ecology* 76:105-134.
- Foster, D.R. and E.R. Boose. 1992. Patterns of forest damage resulting from catastrophic wind in central New England, U.S.A. *Journal of Ecology* 80:79-98.
- Foster, D.R., J.D. Aber, J.M. Melillo, R.D. Bowden, and F.A. Bazzaz. 1997. Forest response to disturbance and anthropogenic stress. *BioScience*, 47:7, p 437.
- Foster, N.W., Nicolson, J.A., and P.W. Hazlett. 1989. Temporal variation in nitrate and nutrient cations in drainage waters from a deciduous forest. *Journal of Environmental Quality* 18: 238-244.
- France, R. L. 1997. The importance of beaver lodges in structuring littoral communities in boreal headwater lakes. *Can. J. Zool.* 75:1009-1013.
- Franklin, J.F.; Spies, T.A.; Pelt, R.V.; Carey, A.B.; Thornburgh, D.A.; Berg, D.R.; Lindenmayer, D.B.; Harmon, M.E.; Keeton, W.S.; Shaw, D.C.; Bible, K.; Chen, J. 2002. Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and Management*. 155: 399-423..
- Franzmann, A.W., and C.C. Schwartz. 1997. Ecology and management of the North American moose. Smithsonian Institution Press and Wildlife Management Institute. 733 pp.
- Freeman, S. 1987. Acid rain trimming health of forests. *The Springfield Union*. 22 May 1987: 17.
- Frelich, L.E. 2002. Forest dynamics and disturbance regimes: studies from temperate evergreen-deciduous forests. *Cambridge Studies in Ecology*. University Press, Cambridge, U.K. 266 pp.
- French, J.M. 1988. MDC-DFWELE land acquisition program: MDC watersheds. Unpublished MDC report.
- French, J.M. and G.S. Buzzell. 1992. Wachusett Reservoir Arborvitae Hedge Report. MDC.
- Garrett, J.D., T. Cassidy, K. McGarigal, K.B. Searcy, and R. Harrington. 2000. Rare, unique, and exemplary natural communities of Quabbin watershed. University of Massachusetts, Department of Natural Resources Conservation report prepared for MDC/DWM.
- Garvey, E. 2000. Natural organic matter fate and transport in the Quabbin Reservoir. PhD Dissertation (unpublished). University of Massachusetts, Amherst, MA. Department of Civil and Environmental Engineering.
- Geldreich, E.E. 1972. Water-borne pathogens. Chapter 9 In: Mitchell (ed.). *Water pollution microbiology*. John Wiley & Sons.
- Gold, A. 1994. Riparian zone hydrology: ramifications for ground water pollution removal. *Proc. Riparian Buffer Zones Conference*. Nov. 21 and 22, Sturbridge, MA. P. Veneman, ed.
- Goldstein, R.A., Chen, C.W., and S.A. Gherini. 1985. Integrated lake-watershed acidification study: summary. *Water, Air, and Soil Pollution* 26:327-337.
- Goodale, C.L., J.D. Aber, and W.H. McDowell. 2000. The long-term effects of disturbance on organic and inorganic nitrogen export in the White Mountains, New Hampshire. *Ecosystems* 3:433-450.
- Gould, D.J. and M.R. Fletcher. 1978. Gull droppings and their effects on water quality. *Water Res.* 12:665-672.
- Gresham, C.A. 1996. Changes in bald cypress-swamp tupelo wetland soil chemistry caused by Hurricane Hugo induced saltwater inundation. In: Haymond, J.L. and W.R. Harms, eds. 1996. *Hurricane Hugo: South Carolina forest land research and management related to the storm*. Gen. Tech. Rep. SRS-5. Asheville, N.C.: U.S. Department of Agriculture, Forest Service, Southern Research Station. Pp. 266-270.

- Hägglund, Å., and G. Sjöberg. 1999. Effects of beaver dams on the fish fauna of forest streams. *For. Ecol. Manage.* 115:259-266.
- Hakanson, L., T. Andersson, and A. Nilsson. 1990. New method of quantitatively describing drainage areas. *Environ. Geol. Sci* 15(1):61-69.
- Hall, C.A.S. and J.W. Day, Jr. 1977. *Ecosystem modeling in theory and practice.* Wiley-Interscience, New York, N.Y. 684p.
- Hammerson, G.A. 1994. Beaver (*Castor canadensis*): Ecosystem alterations, management, and monitoring. *Nat. Areas J.* 14(1):44-57.
- Harr, R.D. and R.L. Fredriksen. 1988. Water quality after logging small watersheds within the Bull Run watershed, Oregon. *Water Resources Bulletin* 24(5): 1103-1111.
- Hartley, B.A. 1975. Current management practices on the Baltimore municipal watersheds. USDA Forest Service, General Technical Report NE-13
- Hartman, E.M., P.E. Austin, and K.M. Keohane. 1992. Quabbin tributaries: water quality data report. Commonwealth of Massachusetts, Department of Environmental Protection and Metropolitan District Commission. Pub. No. 17158-195-50. 196 pp.
- Haymond, J.L. and W.R. Harms, eds. 1996. Hurricane Hugo: South Carolina forest land research and management related to the storm. General Technical Report SRS-5. Asheville, N.C.: U.S. Department of Agriculture, Forest Service, Southern Research Station. 540 pgs.
- Healy, W.M. 1997a. Influence of deer on the structure and composition of oak forests in central Massachusetts. *In: The Science of Overabundance: deer ecology and population management.* Smithsonian Institution Press. 402pp.
- Healy, W.M. 1997b. Thinning New England oak stands to enhance acorn production. *N. Journ. Appl. For.* 14(3):152-156.
- Healy, W.M., R.T. Brooks and P.J. Lyons. 1987. Deer and forests on Boston's municipal watershed after 50 years as a wildlife sanctuary. *IN: Proceedings: deer, forestry, and agriculture: interactions and strategies for management.* June 15-17, 1987. Warren, PA. Allegheny Society of American Foresters.
- Healy, W.M., R.T. Brooks and R.M. DeGraaf. 1989. Cavity trees in sawtimber-size oak stands in central Massachusetts. *North. J. Appl. For.* 6(2):61-65.
- Hedin, L.O., J.J. Armesto and A.H. Johnson. 1995. Patterns of nutrient loss from unpolluted, old-growth temperate forests: evaluation of biogeochemical theory. *Ecology* 76(2), 493-509.
- Hemond, H., K. Eshleman, K. O'Connor, B. Gordesky, and D. Chen. 1987. Surface hydrology and wetland roles in acid deposition on the Bickford Pond watershed. Commonwealth of Massachusetts, Dept. of Environmental Quality Engineering, Project 84-26-WPC. 71 pp.
- Hemond, H.F. and K.N. Eshleman. 1984. Neutralization of acid deposition by nitrate retention at Bickford watershed, Massachusetts. *Water Resources Research.* 20(11):1718-1724.
- Herkert, J.R., R. E. Szafoni, V. M. Kleen, and J.E. Schwegman. 1993. Habitat establishment, enhancement and management for forest and grassland birds in Illinois. *Ill. Dept. Conserv. Div. Nat. Herit.* 1. 19pp.
- Hewlett, J.D. 1978. Forest water quality: an experiment in harvesting and regenerating of Piedmont forest. University of Georgia, Athens, GA.
- Hewlett, J.D. 1982. *Principals of forest hydrology.* University of Georgia Press. Athens, GA. 183 pp.
- Hewlett, J.D. and W.L. Nutter. 1969. *An outline of forest hydrology.* University of Georgia Press, Athens, GA, 137 pp.

- Hibbert, A.R. 1967. Forest treatment effects on waters yield. In: International Symposium on Forest Hydrology, W.E. Sopper and H.W. Lull, eds, Pergamon Press, Oxford, England, pp 527-543.
- Hill, E.P. 1982. Beaver. pp. 256-281 IN: J.A. Chapman and G.A. Feldhammer (eds), Wild mammals of North America: biology, management and economics. John Hopkins Univ. Press.
- Hobbie, J.E. and G.E. Likens. 1973. Output of phosphorus, dissolved organic carbon, and fine particulate carbon from Hubbard Brook watersheds. *Limnology and Oceanography*, 18(5):734-742.
- Hobson, S.S., J.S. Barclay, and S.H. Broderick. 1993. Enhancing wildlife habitats: a practical guide for forest landowners. NE Reg. Agric. Engin. Serv. 172pp.
- Hodkinson, I.D. 1975. Energy flow and organic matter decomposition in an abandoned beaver pond ecosystem. *Oecologia (Berl.)* 21:131-139.
- Hoekstra, T.W., and J. Capp. 1988. Integrating forest management for wildlife and fish. U.S.D.A. For. Serv. Gen. Tech. Rep. NC-122. 63pp.
- Hooper, M.C., K. Aril, and M.J. Lechowicz. 2001. Impact of a major ice storm on an old-growth hardwood forest. *Can. J. Bot.* 79(1): 70-75.
- Hornbeck, J.W. and C.A. Federer. 1975. Effects of management practices on water quality and quantity: Hubbard Brook Experimental Forest, New Hampshire. In: Municipal Watershed Management: Symposium Proceedings. USDA Forest Service General Technical Report NE-13: 58-65.
- Hornbeck, J.W. and J.N. Kochenderfer. 2004. A century of lessons about water resources in Northeastern forests., Chapter 2 In A century of forest and wildland watershed lessons, G.G. Ice and J. D. Stednick, eds., Society of American Foresters, 287 p.
- Hornbeck, J.W., and W.B. Leak. 1992. Ecology and management of northern hardwood forests in New England. U.S.D.A. For. Serv. Gen. Tech. Rep. NE-159. 44p.
- Hornbeck, J.W., C.W. Martin, and C.T. Smith. 1986. Protecting forest streams during whole-tree harvesting. *Northern Journal of Applied Forestry* Vol 3, Sept, pp 97-100.
- Hornbeck, J.W., C.W. Martin, R.S. Pierce, F.H. Bormann, G.E. Likens and J.S. Eaton. 1987. Northern hardwood forest ecosystem: ten years of recovery from clearcutting. U.S.D.A. Forest Service, Northeastern Forest Experiment Station. Broomall, PA. NE-RP-596. 30 p.
- Hornbeck, J.W., E.S. Verry, and C.A. Dolloff. 1994. Managing riparian areas in the northeastern forests: a quest for desired future conditions. Proceedings: Managing forests to meet peoples' needs. Society of American Foresters/Canadian Institute of Forestry Convention, September 18-22, Anchorage, Alaska, 299-304.
- Hornbeck, J.W., G.E. Likens, R.S. Pierce, and F.H. Bormann. 1975. Strip cutting as a means of protecting site and streamflow quality when clearcutting northern hardwoods. In "Forest Soils and Land Management" (B. Bernier and C.F. Winget, eds.), pp.209-225. Presses Univ. Laval, Quebec.
- Hornbeck, J.W., M.B. Adams, E.S. Corbett, E.S. Verry, and J.A. Lynch. 1993. Long-term impacts of forest treatments on water yield: a summary for northeastern USA. *Journal of Hydrology*, 150:323-344.
- Hornbeck, J.W., R.S. Pierce, G.E. Likens, and C.W. Martin. 1975. Moderating the impact of contemporary forest cutting on hydrologic and nutrient cycles. In: The Hydrological Characteristics of River Basins Symposium, IAHS-AISH Publication No. 117, pp 423-433.
- Horsley, S.B. and D.A. Marquis. 1982. Interference by weeds and deer with Allegheny hardwood reproduction. *Can. J. For. Res.* 13:61-69.

- Houghton, J.L., ed. 1996. Climate change 1995: the science of climate change. Inter-Governmental Panel on Climate Change. Cambridge: Cambridge University Press. 584 p.
- Hunt, F.M. 1961. Forest resources on Metropolitan District Commission lands surrounding Quabbin Reservoir. Master's Thesis. University of Massachusetts, Department Forestry and Wildlife Management, Amherst, MA. 128 pp.
- Hunt, F.M. and D.L. Mader. 1970. Low density management - a means to increase timber yields while using less soil moisture. Mass. Agric. Exp. Sta. Bulletin No. 588, November.
- Hunt, P., and D. De Luca. 1997. Survey and management recommendations for grassland and shrubland birds in New Hampshire. Audubon Soc. of NH. 46pp.
- Hunter, M.L. 1990. Wildlife, forests, and forestry; principles of managing forests for biological diversity. Prentice Hall, Englewood Cliffs, N.J. 370 pp.
- Hunter, M.L. Jr. 1999. Maintaining biodiversity in forest ecosystems. Cambridge University Press. 698pp.
- Hunter, W.C., D.A. Beuhler, R.A. Canterbury, J.L. Confer, and P.B. Hamel. 2001. Conservation of disturbance-dependent birds in eastern North America. Wildl. Soc. Bull. 29(2):440-455.
- Hussong, D., J.M. Damare, R.J. Limpert, W.J.L. Sladenk, R.M. Weiner and R.R. Colwell. 1979. Microbial impact of Canada geese (*Branta canadensis*) and whistling swans (*Cygnus columbianus columbianus*) on aquatic ecosystems. Appl. Envir. Microbiol. 37:14-20.
- Ice, G.G. and J.D. Stednick (eds). 2004. A century of forest and wildland watershed lessons. Society of American Foresters, Bethesda, MD. 287 pp.
- Indiana Department of Natural Resources. Undated. Managing woodlands for wildlife. DNR. Manage. Series No. 3. 9pp.
- Iverson, L.R., A.M. Prasad, B.J. Hale, and E.K. Sutherland. 1999. An atlas of current and potential future distributions of common trees of the Eastern United States. Gen. Tech. Rep. NE-265. Radnor, PA. U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 245 p.
- Jensen, P. G., P. D. Curtis, and D. L. Hamelin. 1999. Managing nuisance beavers along roadsides: a guide for highway departments. Cornell University. 14pp.
- Johnson, D.W. and R.I. VanHook, eds. 1989. Analysis of biogeochemical cycling processes in Walker Branch watershed. Springer-Verlag, New York, NY. 401 pp.
- Johnson, E. and M. Stachiw. 1985. Topographic overview. In: Historic and archaeological resources of central Massachusetts, pp 7-20. Massachusetts Historical Commission, Boston.
- Johnson, E.S. and T.F. Mahlstedt. 1984. Prehistoric collections in Massachusetts: The Johnson Craig collection, New Braintree, Massachusetts. Massachusetts Historical Commission, Boston.
- Johnston, C.A. and R. J. Naiman. 1990. Aquatic patch creation in relation to beaver population trend. Ecology. 71(4):1617-1621.
- Johnston, C.A. and R.J. Naiman. 1990. Browse selection by beaver: effects on riparian forest composition. Can. J. For. 20:1036-1043.
- Johnston, W.S., G.K. Maclachlan and G.F. Hopkins. 1979. The possible involvement of seagulls (*Larus* sp.) in the transmission of salmonella in dairy cattle. Veterinary Record. 105:526-527.
- Jones, A.L., and P.D. Vickery. 1998. Conserving grassland birds: managing agricultural lands including hayfields, crop fields, and pastures for grassland birds. Mass. Audubon Society. 17pp.
- Jones, C., C. MacConnell, K. Coleman, P. Cox, P. Falloon, D. Jenkinson, and D. Powlson. 2005. Global climate change and soil carbon stocks; predictions from two contrasting models for the turnover of organic carbon in soil. Global Change Biology. 11: 154-166.

- Jones, C.G., J.H. Lawton, and M. Shachak. 1997. Positive and negative effects of organisms as physical ecosystem engineers. *Ecology*. 78(7):1946-1957.
- Jones, F., P. Smith and D.C. Watson. 1978. Pollution of a water supply catchment by breeding gulls and the potential environmental health implications. *J. Inst. Wat. Eng. & Sci.* 32:469-482.
- Jones, W.R. 1951. The deer herd of Prescott Peninsula, Quabbin Reservation, and its ecology. M.S. Thesis, University of Massachusetts, Amherst. 126 pp.
- Jordan, C.F. 1982. Amazon rain forests. *American Scientist* 70:394-401.
- Kelty, M.J., D.B. Kittredge, Jr., T. Kyker-Snowman, and A.D. Leighton. 2003. The conversion of even-aged stands to uneven-aged structure in Southern New England. *Northern Journal of Applied Forestry* 20(3), pp 109-116.
- Kimple, A. and W. Schuster. 2002. Spatial patterns of HWA damage and impacts on tree physiology and water use in the Black Rock Forest, southern New York. In: *Proceedings: hemlock woolly adelgid in the eastern United States*. February 5-7, 2002. B. Onken (ed). Rutgers – the State University of New Jersey, East Brunswick, NJ. 403 p.
- King, D.A. 1986. Tree form, height growth, and susceptibility to wind damage in *Acer saccharum*. *Ecology*, 67: 980-990.
- King, S.L., B.D. Keeland, and J.L. Moore. 1998. Beaver lodge distributions and damage assessments in a forested wetland ecosystem in the southern United States. *For. Ecol. Manage.* 108:1-7.
- Kizlinski, M.L., D.A. Orwig, R.C. Cobb, and D.R. Foster. 2002. Direct and indirect ecosystem consequences of an invasive pest on forests dominated by eastern hemlock. *Journal of Biogeography*, 29:1489-1503.
- Klein, R.M. and T.D. Perkins. 1988. Primary and secondary causes and consequences of contemporary forest decline. *The Botanical Review*. 54(1), Jan-Mar, 43 pp.
- Klotz, R. L. 1998. Influence of beaver ponds on the phosphorus concentration of streamwater. *Can. J. Fish. Aquat. Sci.* 55:1228-1235.
- Kochenderfer, J.N. and G.M. Aubertin. 1975. Effects of management practices on water quality and quantity: Fernow Experimental Forest, West Virginia. In: *Municipal Watershed Management Symposium Proceedings*, USDA Forest Service, General Technical Report, NE-13, pp 14-24.
- Kochenderfer, J.N. and J.D. Helvey. 1974. Gravel greatly reduces soil losses from minimum-standard forest roads. *USDA Forest Service, Parsons, West Virginia*.
- Kochenderfer, J.N., P.J. Edwards, and F. Wood. 1997. Hydrological impacts of logging an Appalachian watershed using West Virginia's best management practices. *Northern Journal of Applied Forestry* 14(4): 207-218.
- Krug, E.C. and C.R. Frink. 1983. Acid rain on acid soil: a new perspective. *Science*. 221:520-525.
- Kyker-Snowman, T.D. 1989a. Quabbin forest regeneration study. MDC Publication.
- Kyker-Snowman, T. 1989b. Logging systems evaluation on the Wachusett Reservoir, November, 1989: summary of evaluation and analysis. MDC Publication
- LaDeau, S.L. and J.E. Clark. 2001. Rising CO₂ levels and the fecundity of forest trees. *Science*. 292:95-98.
- Larsen, D.R. and P.S. Johnson. 1998. Linking the ecology of natural oak regeneration to silviculture. *Forest Ecology and Management*, Vol. 106, no. 1, pp. 1-7.
- Larson, F.R., P.F. Ffolliott, W.O. Rasmussen, and D.R. Carder. 1979. Estimating impacts of silvicultural management practices on forest ecosystems. In: *Best Management Practices for Agriculture and*

- Silviculture: Proceedings of the 1978 Cornell Agricultural Waste Management Conference, Ann Arbor Science Publishers Inc., Ann Arbor, Michigan.
- Lawrence, G.B., D.A. Burns and P.S. Murdoch. 1993. Work plan of the Neversink watershed study. United States Geological Survey. Albany, N.Y.
- Leak, W.B. and C.W. Martin. 1975. Relationship of stand age to streamwater nitrate in New Hampshire. USDA Forest Service Research Note NE-211, 5 pp.
- Leak, W.B. and J.H. Gottsacker. 1985. New approaches to uneven-age management in New England. Northern Journal of Applied Forestry. 2:28-31.
- Leonard, W.C., R.F. Yuretich, and S.J. Pohanka. 1984. Hydrogeology and processes of acid neutralization in Cadwell Creek, Quabbin Reservation, Massachusetts. Water Resources Research Center, University of Massachusetts, Pub. No. 148. 122 pp.
- Levno, A. and J. Rothacher. 1969. Increases in maximum stream temperatures after slash burning in a small experimental watershed. Res. Note PNW-110. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. 7p.
- Lewis, G.P. and G.E. Likens. 2000. Low stream nitrate concentrations associated with oak forests on the Allegheny High Plateau of Pennsylvania. Water Resources Res 36:3091-3094.
- Li, R.M., K.G. Eggert, and D.B. Simons. 1979. Simulation of stormwater runoff and sediment yield for assessing the impact of silvicultural practices. In: Best Management Practices for Agriculture and Silviculture: Proceedings of the 1978 Cornell Agricultural Waste Management Conference, Ann Arbor Science, Ann Arbor, Michigan.
- Litvaitis, J.A. 2001. Importance of early successional habitats to mammals in eastern forests. Wild. Soc. Bull. 29(2):466-473.
- Lorimer, C. 1977. The presettlement forest and natural disturbance cycle of northeastern Maine. Ecology. 58(1):139-148.
- Lovett, G.M., K.C. Weathers, and M.A. Arthur. 2002. Control of nitrogen loss from forested watersheds by soil carbon:nitrogen ratio and tree species composition. Ecosystems 5: 712-718.
- Lovett, G.M., K.C. Weathers, and W.V. Sobczak. 2000. Nitrogen saturation and retention in forested watersheds of the Catskill Mountains, New York. Ecological Applications 10(1):73-84.
- Lovett, G.M. and H. Rueth. 1999. Soil nitrogen transformations in beech and maple stands along a nitrogen deposition gradient. Ecol Appl 9:1330-1344.
- Lowrance, R. 1994. Technical overview of water quality functions of riparian forest buffer systems. Proc. Riparian Buffer Zones Conference. Nov. 21 and 22, Sturbridge, MA. P. Veneman ed.
- Lowrance, R. and G. Vellidis. 1994. A conceptual model for assessing ecological risk to water quality function of bottomland hardwood forests. Environmental Management 19(2):239-258.
- Lull, H.W. and K.G. Reinhart. 1967. Increasing water yield in the northeast by management of forested watersheds. USDA Forest Service Research Paper NE-66.
- Lull, H.W. and K.G. Reinhart. 1972. Forests and floods in the eastern United States. In: Readings in forest hydrology, Eschner, A.R. and P.E. Black, Eds, MSS Information Corp., New York, N.Y.
- Lynch, J.A., W.E. Sopper, E.S. Corbett, and D.W. Aurand. 1975. Effects of management practices on water quality and quantity: the Penn State Experimental Watersheds. In: Municipal Watershed Management Symposium Proceedings, USDA Forest Service, General Technical Report, NE-13, pp 32-46.
- Mackay, D.S. and L.E. Band. 1997. Forest ecosystem processes at the watershed scale: dynamic coupling of distributed hydrology and canopy growth. Hydrological Processes, 11:1197-1217.

- Mader, D.L., W.P. MacConnell, and J.W. Bauder. 1972. The effect of riparian vegetation control and stand density reduction on soil moisture in the riparian zone. University of Massachusetts – Amherst, College of Food and Natural Resources, Agricultural Experiment Station. Research Bulletin 597.
- Magill, A. and J. Aber. 1993. Changes in nitrogen mineralization, nitrification and nitrogen leaching in the chronic N plots at Harvard Forest. IN: Long Term Ecological Research at Harvard Forest, Foster, D.R. and D.R. Smith eds., 109 pp.
- Malben, G. and J.B. Foote. 1955. Beaver activity and water quality. *J. Amer. Water Works Assoc.* 47:503-507.
- Marchand, P.J. 1987. North woods: an inside look at the nature of forests in the Northeast. Appalachian Mountain Club, Boston, MA.
- Maret, T.J., M. Parker, and T.E. Fannin. 1987. The effect of beaver ponds on the nonpoint source water quality of a stream in southwestern Wyoming. *Wat. Res.* 21(3):263-268.
- Margolis, B.E., M.S. Castro, and R.L. Raesly. 2001. The impact of beaver impoundments on the water chemistry of two Appalachian streams. *Can. J. Fish. Aquat. Sci.* 58: 2271-2283.
- Marks, P.L. and F.H. Bormann. 1972. Revegetation following forest cutting: mechanisms for return to steady-state nutrient cycling. *Science.* 176:914-915.
- Marquis, D.A., P.L. Eckert, and P.A. Roach. 1976. Acorn weevils, rodents, and deer all contribute to oak regeneration difficulties in Pennsylvania. USDA, For. Service Research Paper NE-356, 5 pp.
- Marquis, D.A., T.J. Grisez, J.C. Bjorkbom, and B.A. Roach. 1975. Interim guide to regeneration of Allegheny hardwoods. USDA Forest Service Gen. Tech. Rep. NE-19.
- Marquis, David A. 1991. Uneven-Aged Management in the Cherry-Maple Forest. Proceedings of the Allegheny Society of American Foresters Symposium on Uneven-aged Management.
- Martin, C.W. 1988. Soil disturbance by logging in New England - review and management recommendations. *Northern Journal of Applied Forestry* Vol 5, March, pp 30-34.
- Martin, C.W. and R.S. Pierce. 1980. Clearcutting patterns affect nitrate and calcium in streams in New Hampshire. *Journal of Forestry.* May: 268-272.
- Martin, C.W., D.S. Noel, and L.A. Fedrer. 1984. Effects of forest clearcutting in New England on stream chemistry. *Journal of Environmental Quality* Vol 13, pp 204-210.
- Maser, C., S.P. Cline, K. Cromack Jr., J.M. Trappe, and E. Hansen. 1988. What we know about large trees that fall to the forest floor. Chapter 2, in: *From the Forest to the Sea: A Story of Fallen Trees.* USDA Forest Service, General Technical Report, PNW-GTR-229.
- Mather, M. E. 1993. The Quabbin ecosystem: an interdisciplinary ecological review. Final report of the seminar. University of Massachusetts, Department of Forestry and Wildlife Management.
- Matthews, J.D. 1989. *Silvicultural systems.* Oxford University Press, New York, NY.
- Mayer, P.M., S.K. Reynolds, M.D. McCutchen, and T.J. Canfield. 2006. Riparian buffer width, vegetative cover, and nitrogen removal effectiveness: a review of current science and regulations. EPA/600/R-05/118. Cincinnati, OH, U.S. Environmental Protection Agency.
- McAvoy, D.C. 1989. Episodic response of aluminum chemistry in an acid-sensitive Massachusetts catchment. *Water Resources Research.* 25(2):233-240.
- McCull, J.G. and D.F. Grigal. 1975. Forest fire: effects on phosphorus movement to lakes. *Science.* 188:1109-1111.

- McEvoy, T.J. 2004. Positive impact forestry: a sustainable approach to managing woodlands. Island Press, Washington, D.C. 268 pp.
- McInnes, P.F., R.J. Naiman, J.Pastor, and Y.Cohen. 1992. Effects of moose browsing on vegetation and litter of the boreal forest, Isle Royale, Michigan, USA. *Ecology*. 73(6): 2059-2075.
- McKee, W. 1993. Personal communication. Southeastern Forest Experiment Station, U.S.D.A. Forest Service.
- McLaren, B.E., S.P. Mahoney, T.S. Porter, and S.M.Oosenbrug. 2000. Spatial and temporal patterns of use by moose of pre-commercially thinned, naturally-regenerating stands of balsam fir in central Newfoundland. *Forest Ecology and Management*. 113: 179-196.
- McManus, K.A., K.S. Shields, and D.R. Souto (eds.). 2000. Proceedings: symposium on sustainable management of hemlock ecosystems in eastern North America: June 22-24, 1999. Durham, N.H. Gen. Tech. Rep 267. Newtown Square, PA: USDA Forest Service, Northeastern Research Station. 237 p. .
- McRae, G., and C. J. Edwards. 1994. Thermal characteristics of Wisconsin headwater streams occupied by beaver: implications for brook trout habitat. *Trans. Am. Fish. Soc.* 123:641-656.
- McShea, W.J., and J.H. Rappole. 1997. Herbivores and the ecology of forest understory birds. *In: The Science of Overabundance: deer ecology and population management*. Smithsonian Institution Press. 402pp.
- McShea, W.J., H.B. Underwood, and J.H. Rappole. 1997. *The science of overabundance: deer ecology and population management*. Smithsonian Institution Press. 402 pp.
- MDC, Division of Watershed Management. 1988. Water quality report: Quabbin Reservoir and Wachusett Reservoir: watershed monitoring results. Unpublished MDC report.
- MDC, Division of Watershed Management. 1991. Quabbin Reservation white-tailed deer impact management plan. Unpublished MDC report.
- MDC, Division of Watershed Management. 2002. Sudbury Reservoir Watershed System Public Access Plan Update.
- MDC/MWRA. 1991. Watershed protection plan: Quabbin Reservoir and Ware River watersheds. Prepared by Rizzo Assoc. and CH2M Hill. February 28, 1991.
- Meentemeyer, R.K. and D.R. Butler. 1999. Hydrogeomorphic effects of beaver dams in Glacier National Park, Montana. *Phys. Geo.* 20 (5):436-446.
- Melillo, J.M., J.D. Aber, P.A. Steudler and J.P. Schimel. 1983. Denitrification potentials in a successional sequence of northern hardwood forest stands. *Environmental Biogeochemistry Ecological Bulletin, Stockholm*. 35:217-228.
- Metts, B. S., J. D. Lanham, and K. R. Russell. 2001. Evaluation of herpetofaunal communities on upland streams and beaver-impounded streams in the upper piedmont of South Carolina. *Amer. Midl. Nat.* 145.
- Metzmaker, J.B. and A.E. Rosquist. 1986. Watershed monitoring to assess the risk of a *Giardia lamblia* contamination of unfiltered municipal water supplies. *International Symposium on Water-Related Health Issues, American Water Resources Association, November*, pp 145-151.
- Meyer, J.L. and G.E. Likens. 1979. Transport and transformation of phosphorus in a forest stream ecosystem. *Ecology*. 60(6):1255-1269.
- Micks, P., R. Boone and S. Scott. 1994. Abiotic N fixation by forest floor material from the chronic N plots. *Abstr. 5th Annual Harvard Forest Ecology Symposium*. April 25, 1994.

- Miller, D.R., M.J. Focazio, and M.A. Dickinson. 1986. A user's guide to a model for estimating the hydrological effects of land use change (BROOK). Cooperative Extension, University of Massachusetts, University of Connecticut.
- Mladenoff, D.J. 1987. Dynamics of nitrogen mineralization and nitrification in hemlock and hardwood treefall gaps. *Ecology*. 68:1171-1180.
- Mladenoff, D.J., M.A. White, J. Pastor, and T.R. Crow, 1993. Comparing spatial pattern in unaltered old-growth and disturbed forest landscapes. *Ecological Applications* 3(2):294-306.
- Monaghan, P., C.B. Shedden, K. Ensor, C.R. Fricker and R.W.A. Girdwood. 1985. Salmonella carriage by herring gulls in the Clyde area of Scotland in relation to their feeding ecology. *J. Appl. Ecol.* 22:669-680.
- Monzingo, D.L., Jr. and C.P. Hibler. 1987. Prevalence of *Giardia* sp. in a beaver colony and the resulting environmental contamination. *J. Wildlf. Dis.* 23(4):576-585.
- Moore, S.T. 1966. Management of a municipal supply watershed. *Symposium of Forest Watershed Management: Practical Aspects*. Society of American Foresters, Corvallis, Oregon. pp 61-68.
- More, M.E. and J.W. Soper. 1990. GEIR forestland management practices. Massachusetts Department of Environmental Management, EOEEA File Number 6307, 475 pp.
- Motzkin, G., D. Foster, A. Allen and J. Harrod. 1994. Vegetation and regional physiographic and disturbance gradients. Abstracts from the 5th Annual Harvard Forest Ecology Symposium. Petersham, M.A.
- Mrazik, B.R, D.L. Mader, and W.P. MacConnell. 1980. Integrated watershed management: an alternative for the northeast. Mass. Agricultural Experiment Station, Research Publication Number 664, Univ of Mass, Amherst.
- Murdoch, P.S. 1993. Personal communication. USGS Water Res. Div., Albany, N.Y.
- Murdoch, P.S., D.A. Burns, and G.B. Lawrence. 1998. Relation of climate change to the acidification of surface waters by nitrogen deposition. *Environmental Science and Technology*. 32:11, 1642-1647.
- Murdoch, P.S. and J.L. Stoddard. 1992. The role of nitrate in the acidification of streams in the Catskill Mountains of New York. *Water Resources Research*. 28(10):2707-2720.
- MWRA/MDC. 1997. Watershed Protection Plan: Sudbury Watershed and Framingham Reservoir No.3. Prepared with assistance of Comprehensive Environmental, Inc., 95 pp + app.
- Nadelhoffer, K.J., J.D. Aber and J.M. Melillo. 1984. Seasonal patterns of ammonium and nitrate uptake in nine temperate forest ecosystems. *Plant and Soil*, 80: 321-335.
- Naiman, R.J. and J.M. Melillo. 1984. Nitrogen budget of a subarctic stream altered by beaver (*Castor canadensis*). *Oecologia (Berlin)* 62:150-155.
- Naiman, R.J., C.A. Johnston and J.C. Kelly. 1988. Alteration of North American streams by beaver. *BioScience* 38(11):753-762.
- Naiman, R.J., G. Pinay, C.A. Johnston, and J. Pastor. 1994. Beaver influences on the long-term biogeochemical characteristics of boreal forest drainage networks. *Ecology* 75(4):905-921.
- Naiman, R.J., J.M. Melillo and J.E. Hobbie. 1986. Ecosystem alteration of boreal forest streams by beaver (*Castor canadensis*). *Ecology* 67(5):1254-1269.
- National Research Council. 2000. Watershed management for potable water supply: assessing the New York City strategy. National Academy Press, Washington, D.C. 549 pp.
- Neville, R. 1996. personal communication. USDA Forest Service, Durham, NH.

- NHESP (Massachusetts Natural Heritage and Endangered Species Program). 2006. Massachusetts natural heritage atlas, 12th edition. Printed and distributed by the Massachusetts Division of Fisheries and Wildlife Natural Heritage and Endangered Species Program, Westborough, MA. www.nhesp.org
- Nihlgard, B. 1985. The ammonium hypothesis - an additional explanation to the forest dieback in Europe. *Ambio*. 14(1):2-8.
- Nilsson, A. and L. Hakanson. 1992. Relationships between drainage area characteristics and lake water quality. *Environ. Geol. Water Sci.* 19(2):75-81.
- Nolet, B.A., A. Hoekstra, and M.M. Ottenheim. 1994. Selective foraging on woody species by the beaver (*Castor fiber*), and its impact on a riparian willow. *Forest. Biol. Cons.* 70:117-128.
- Norton, D.A. 1999. Chapter 16, Forest Reserves in M.L. Hunter, Jr. 1999. Maintaining biodiversity in forest ecosystems. Cambridge University Press. 698pp.
- Norvell, W.A. and C.R. Frink. 1975. Estimation of potential nutrient enrichment of Lake Wononscopomuc by wild geese. unpubl. manuscript. Conn. Agric. Exp. Station., New Haven.
- Novak, M. 1987. In: M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild furbearer management and conservation in North America. Ontario Ministry of Natural Resources, Toronto. Pp. 283-312
- Novotny, V. 2003. Water quality: diffuse pollution and watershed management. John Wiley & Sons, Inc. NY, NY. 864 p.
- O'Connor, R.W. 1982. Preventing soil erosion on municipal watersheds. Unpublished paper. S.U.N.Y., Syracuse, NY.
- O'Connor, R. and J. Zilligen. 1994. Public access plan for the MDC Sudbury watershed, MDC document, 158 pp.
- Odum, E. P. 1969. The strategy of ecosystem development. *Science*. 164:262-270.
- O'Hara, K.L. 1998. Silviculture for structural diversity: a new look at multiaged systems. *Journal of Forestry*. Volume 96, Number 7, pp. 4-10.
- Ollinger, S., J. Aber and C. Federer. 1994. A GIS modeling approach to estimating regional carbon and water balances for the Northeastern U.S. Abstracts from the 5th Annual Harvard Forest Ecology Symposium. Petersham, MA.
- Onken, B. (ed.) 2002. Proceedings: hemlock woolly adelgid in the eastern United States.
- Orwig, D.A., D.R. Foster, D.L. Mausell. 2002. Landscape patterns of hemlock decline in New England due to the introduced hemlock woolly adelgid. *Journal of Biogeography*, 29: 1475-1487.
- Orwig, D.A. and D.B. Kittredge. 2005. Silvicultural options for managing hemlock forests threatened by hemlock woolly adelgid. IN: U.S.D.A. Forest Service, 2005. B. Onken and R. Reardon, Compilers. Third symposium on hemlock woolly adelgid in the eastern United States. Asheville, North Carolina. February 1-3, 2005. Online: http://www.na.fs.fed.us/fhp/hwa/pub/2005_proceedings/index.shtml
- Ottenheimer, D.G. 1992. Hurricane susceptibility and water quality at Quabbin Forest, Massachusetts. M.S. Thesis. State University of New York, Syracuse. 70pp, +app.
- OWM. 2004. Hemlock woolly adelgid (*Adelges tsugae* Annand) on DWSP lands: problem extent and proposed management approach. Unpublished agency report.
- Parker, D. 1988. Forest endangered: acid rain, ozone damage worsening. *Springfield Union*. 16 June 1988.

- Parsons, W.F.J., D.H. Knight and S.L. Miller. 1994. Root gap dynamics in lodgepole pine forest: nitrogen transformations in gaps of different size. *Ecological Applications*. 4(2):354-362.
- Paterson, K.G. and J.L. Schnoor. 1992. Fate of alachlor and atrazine in a riparian zone field site. *Water Environment Research*. 64(3):274-283.
- Patric, J.H. 1976. Soil Erosion in the eastern forest. *J. Forestry Oct.*, pp 671-677.
- Patric, J.H. 1977. Soil erosion and its control in eastern woodlands. *Northern Logger and Timber Processor Vol 25, No. 11, May*, pp 4-6.
- Patric, J.H. 1978. Harvesting effects on soil and water in eastern hardwoods forests. *Southern Journal of Applied Forestry* 2(3):66-73.
- Patric, J.H. 1980. Effects of wood products harvest on forest soil and water relations. *Journal of Environmental Quality*, 9(1): (73-80).
- Patric, J.H. and E.M. Gould. 1976. Shifting land use and the effects on river flow in Massachusetts. *Journal of American Water Works Association*. (Jan): 41-45.
- Patric, J.H, J.O. Evans, and J.D. Helvey. 1984. Summary of sediment yield data from forested land in the United States. *J. For.* 82(2): 101-104.
- Payne, N.F. and F.C. Bryant. 1994. *Techniques for wildlife habitat management of uplands*. McGraw-Hill, Inc. 840pp.
- Peek, J.M., and K.I. Morris. 1998. Status of moose in the contiguous United States. *Alces*. 34(2): 423-434.
- Perkey, A.W., B.L. Wilkens, and H.C. Smith. 1993. *Crop Tree Management in Eastern Hardwoods*, U.S. Forest Service Publication: NA-TP-19-93.
- Peters, N.E. and P.S. Murdoch. 1985. Hydrogeologic comparison of an acidic-lake basin with a neutral-lake basin in the west-central Adirondack Mountains, New York. *Water, Air, and Soil Pollution*. 26:387-402.
- Petersen, C. and G.C. Smith. 1989. Are the forests of Massachusetts in decline?. IN: *Current issues in forestry*, Cooperative Extension, University of Mass., Amherst, Vol I, No 2, May, 4 pp.
- Pickett, S.T.A. and P.S. White, eds.. 1985. *The ecology of natural disturbance and patch dynamics*. Academic Press Inc., New York, 472 pp.
- Portnoy, J.W. 1990. Gull contributions of phosphorous and nitrogen to a Cape Cod kettle pond. *Hydrobiologia* 202:61-69.
- Putz, F.E. and R.R. Sharitz. 1991. Hurricane damage to old-growth forest in Congaree Swamp National Monument, South Carolina, U.S.A. *Can. J. For. Res.* 21: 1765-1770.
- Raup, H.M. 1957. Vegetational adjustment to the instability of site. in *Proc. 6th Tech. Meet. 1956 Int. Un. Conserv. Nature Nat. Resources*, Edinburgh. pp 36-48.
- Reese, K.P. and J.D. Hair. 1976. Avian species diversity in relation to beaver pond habitats in the Piedmont region of South Carolina. *Proc. Southeast Assoc. F & W Agencies Conf.* 30:437-447.
- Reich, P.B. and R.G. Amundson. 1985. Ambient levels of ozone reduce net photosynthesis in tree and crop species. *Science*. 230:566-570.
- Reid, J. 1992. Personal communication. Hampshire College, Amherst, MA.
- Reid, W., B.L. Parker, M. Skinner, S. Gouli, and H.B. Teillon. 2002. Insect-killing fungi for management of hemlock woolly adelgid: a review of progress. In: *Proceedings: hemlock woolly adelgid in the eastern United States*. February 5-7, 2002. B. Onken (ed). Rutgers – the State University of New Jersey, East Brunswick, NJ. 403 p.

- Reinhart, K.G. and A.R. Eschner. 1962. Effects on streamflow of four different forest practices in the Allegheny Mountains, J. Geophys. Research. 67:2433-2445.
- Reinhart, R.H. and J.A. Hroncich. 1990. Source water quality management. pp. 189-228 IN: F.W. Pontius (ed). Water quality and treatment: a handbook of community water supplies. AWWA. McGraw-Hill.
- Remillard, M.M., G.K. Gruending and D.J. Bogucki. 1987. Disturbance by beaver (*Castor canadensis*) and increased landscape heterogeneity. Chapt. 6 IN: M.G.Turner (ed), Ecological Studies: Vol 64. Landscape heterogeneity and disturbance. Springer-Verlag, N.Y.
- Reuss, J.O. and D.W. Johnson. 1986. Acid deposition and the acidification of soils and waters. Ecological Studies. Springer-Verlag; New York, N.Y.
- Rittmaster, R.L. and J.B. Shanley. 1990. Effects of atmospheric wet deposition on the water quality of two streams in the Quabbin Reservoir watershed, central Massachusetts, 1983-85. U.S. Geological Survey. Water-Resources Investigations Report 90.
- Rittmaster, R.L. and J.B. Shanley. 1995. Factors affecting water quality and net flux of solutes in two stream basins in the Quabbin Reservoir drainage basin, central Massachusetts, 1983-85. U.S. Geological Survey. Water-Resources Investigations Report 94-4003.
- Roberts, M.R. and F.S. Gilliam. 1995. Patterns and mechanisms of plant diversity in forested ecosystems: implications for forest management. *Ecological Applications*, Vol. 5, No. 4, pp. 969-977
- Robinson, J.B., H.R. Whiteley, W. Stammers, N.K. Kaushik, and P. Sain. 1979. The fate of nitrate in small streams and its management implications. In: Best management practices for agriculture and silviculture: Proceedings of the 1978 Cornell Agricultural Waste Management Conference, Ann Arbor Science Publishers, pp 247-259.
- Rose, J.B. 1988. Occurrence and significance of *Cryptosporidium* in water. J. Amer. Wat. Works Assoc.:53-58.
- Rothacher, J. 1970. Managing forest land for water quality. IN: Proceedings of the Joint FAO/U.S.S.R. international symposium on forest influences and watershed management, Moscow, U.S.S.R. by USDA Forest Service.
- Runkle, J. R. 1985. Disturbance regimes in temperate forests. In: The Ecology of Natural Disturbance and Patch Dynamics. S. T. A. Pickett and P. S. White (editors), Academic Press, Orlando, FL, pp. 17-33.
- Runkle, J.R. 1982. Patterns of disturbance in some old-growth mesic forests of eastern North America. *Ecology* 63(5): 1533-1546
- Russell, E.W.B. 1983. Indian-set fires in the forests of the Northeastern United States. *Ecology*. 64: 78-88.
- Russell, H.S. 1976. A long, deep furrow: three centuries of farming in New England. University Press of New England. Hanover, N.H. 672 pp.
- Russell, K.R., C.E. Moorman, J.K. Edwards, B.S. Metts, and D.C. Guynn, Jr. 1999. Amphibian and reptile communities associated with beaver (*Castor canadensis*) ponds and unimpounded streams in the piedmont of South Carolina. *J. Freshwater Ecol.* 14(2):149-160.
- S.U.N.Y. 1981. New York river basins special studies: a proposal plan of forestland watershed management for Schoharie watershed. Syracuse, N.Y. Unpublished. (cited in O'Connor, 1982).
- Sample, D.W. and M.J. Mossman. 1997. Managing habitat for grassland birds: a guide for Wisconsin. Wis. Dept. Nat. Res. 154pp.
- Satterlund, D.R. and P.W. Adams. 1992. Wildland watershed management. 2nd edition. John Wiley & Sons, Inc. New York, N.Y.

- Savill, P.S. 1983. Silviculture in windy climates. *Forestry Abstracts*. 44(8):473-488.
- Schindler, D.W. 1988. Effects of acid rain on freshwater ecosystems. *Science*. 239:149-157.
- Schindler, D.W., R.W. Newbury, K.G. Beaty, J. Prokopowich, T. Ruszczynski, and J.A. Dalton. 1980. Effects of a windstorm and forest fire on chemical losses from forested watersheds and on the quality of receiving streams. *Canadian Journal of Fisheries Aquatic Science*. 37:328-334.
- Schofield, C.L., J.N.Galloway, and G.R. Hendry. 1985. Surface water chemistry in the ILWAS Basins. Abstract. *Water, Air, and Soil Pollution*. 26:403-423.
- SCS. 2004. Scientific Certification System's "Certification Evaluation Report for the Natural Forests Managed by the Commonwealth of Massachusetts". Online: <http://www.mass.gov/envir/forest/pdf/forestgreencertreport.pdf> . 175 pp.
- Schueler, T. 1987. Controlling urban runoff: a practical manual for planning and designing urban BMPs. Prepared for the Washington Metropolitan Water Resources Planning Board.
- Scudder, J.H. 1949. A preliminary study of forest succession in central New England. Masters thesis, Harvard University. 69 pp.
- Searcy, K.B. 1995. Quabbin rare plant survey, June-August, 1995. Final report to MDC.
- Searcy, K.B. 1996. 1996 rare plant survey: Quabbin, Ware River, Wachusett, and Sudbury watersheds. Final report to MDC.
- Shanley, J.B. 1993. Personal communication. United States Geological Survey.
- Sheehan, K.C. 1982. Development of environmental impact criteria for recreational use of the Quabbin Reservoir. Master's thesis, Dept. Civil Engineering, University of Massachusetts. 95 pp.
- Shepard, J.P., M.J. Mitchell, T.J. Scott, and C.T. Driscoll. 1990. Soil solution chemistry of an Adirondack spodosol: lysimetry and N dynamics. *Canadian Journal of Forest Resources*, 20:818-824.
- Skeffington, R.A. and E.J. Wilson. 1988. Excess nitrogen deposition: issues for consideration. *Environmental Pollution*. 54:159-184.
- SmartWood. 1997. Certification assessment report of the Metropolitan District Commission's Quabbin Reservoir lands, Commonwealth of Massachusetts. National Wildlife Federation, Northeast Natural Resource Center and The Rainforest Alliance's SmartWood Program. June 1997, 60 pp.
- Smith, D.M., B.C. Larson, M.J. Kelty, P.M.S. Ashton. 1997. *The practice of silviculture: applied forest ecology*. John Wiley and Sons, New York, NY. 560 pp.
- Smith, D.M. 1986. *The practice of silviculture*, eighth edition. John Wiley and Sons, New York, NY. 527 pp.
- Smith, M.E., C.T. Driscoll, B.J. Wyskowski, C.M. Brooks and C.C. Cosentini. 1991. Modification of stream ecosystem structure and function by beaver (*Castor canadensis*) in the Adirondack Mountains, New York. *Can. J. Zool.* 69:55-61.
- Smith, W.H. 1981. Air pollution-effects on the structure and function of the temperate forest ecosystem. In: *Air pollution and forests*. Springer-Verlag, New York. 379 pp.
- Snodgrass, J.W. 1997. Temporal and spatial dynamics of beaver-created patches as influenced by management practices in a south-eastern North American landscape. *J. Appl. Ecol.* 34:1043-1056.
- Snodgrass, J.W. and G.K. Meffe. 1998. Influence of beavers on stream fish assemblages: effects of pond age and watershed position. *Ecology*. 79(3):928-942.
- Snow, D.R. 1980. *The archaeology of New England*. Academic Press, New York, N.Y.

- Snyder, E.J., and K. Bontaitis. 2001. Moose fact sheet. University of New Hampshire Cooperative Extension. 4 pp.
- Sorrie, B.A. 1989. Massachusetts flora: a review of current distribution and conservation of rare species. *Rhodora* 91:116-120.
- Speranza, D.E. 1988. An investigation of *Giardia lamblia* in the Quabbin and Wachusett reservoirs. M.S. Thesis, University of Massachusetts, Amherst.
- Spurr, S.H. and B.V. Barnes. 1980. Forest ecology (third edition). John Wiley and Sons, Inc. New York, NY. 687 pp.
- Standridge, J.H., J.J. Delfino, L.B. Kleppe and R. Butler. 1979. Effect of waterfowl (*Anas platyrhynchos*) on indicator bacteria populations in a recreational lake in Madison, Wisconsin. *Appl. Environ. Microbiol.* 38:547-550.
- Stednick, J.D. 2000. Timber management. In: Dissmeyer, G.E., ed. Drinking water from forests and grasslands; A synthesis of the scientific literature. U.S.D.A. Forest Service, General Technical Report SRS-39. Asheville, N.C.: 103-119.
- Stekl, P.J. 1985. A hydrogeologic investigation of two small watersheds of the eastern Quabbin basin. Masters Thesis, University of Massachusetts.
- Stephens, E.P. 1956. The uprooting of trees: a forest process. *Proceedings of the Soil Science Society of America.* 20:113-116.
- Stoddard, J.L. 1991. Trends in Catskill stream water quality: evidence from historical data. *Water Resources Research.* 27(11):2855-2864.
- Stone, E.L. 1973. The impact of timber harvest on soils and water. In: Report of the President's Advisory Panel on Timber and the Environment, April, pp 427-467.
- Stone, E.L., W.T. Swank, and J.W. Hornbeck. 1979. Clearcutting versus alternative timber harvest and regeneration systems: impacts on soils and streamflow in the eastern deciduous region. USDA Forest Service.
- Storey, H.C. and I.C. Reigner. 1970. Vegetation management to increase water yields. In: Proceedings of the Symposium on Interdisciplinary Aspects of Watershed Management, August 3-6, Montana State University, American Society of Civil Engineers, New York, pp 271-293.
- Stuart, G.W. and P.J. Edwards. 2006. Concepts about forests and water. *Northern Journal of Applied Forestry* 23:1, 11-19.
- Suurballe, N.C. 1992. Effects of a wetland on quality of Natty Pond Brook, Massachusetts, 1985-86. U.S. Geological Survey. Water-Resources Investigations Report 91-4144. 52 pp.
- Swank, W.T. 1972. Water balance, interception, and transpiration studies on a watershed in the Puget lowland region of western Washington. Ph.D. Dissertation, University of Washington, Seattle.
- Swank, W.T., J.B. Waide, D.A. Crossley, Jr., and R.L. Todd. 1981. Insect defoliation enhances nitrate export from forest ecosystems. *Oecologia* 51:297-299.
- Tamm, C.O. 1991. Nitrogen in terrestrial ecosystems: questions of productivity, vegetational changes, and ecosystem stability. *Ecological Studies.* Springer-Verlag, New York, N.Y. 116p.
- Thompson, C. H. and T.D. Kyker-Snowman. 1989. Evaluation of non-point source pollution problems from crossing streams with logging equipment and off-road vehicles in Massachusetts: 1987-1988. Report produced as part of Generic Environmental Impact Report on Forest Management in Massachusetts. Massachusetts Department of Environmental Management.
- Thompson D.Q. and R.H. Smith 1970. The forest primeval in the Northeast--a great myth? In: Proc Ann Tall Timbers Fire Ecology Conf 10: 255-265, R. Komarek, ed.. Tallahassee, FL.

- Thompson, F.R., and R.M. DeGraaf. 2001. Conservation approaches for woody, early successional communities in the eastern United States. *Wild. Soc. Bull.* 29(2):483-494.
- Thompson, F.R. III and D.R. Dessecker. 1997. Management of early successional communities in central hardwood forests. USDA For. Serv. Gen. Tech. Rep. NC-195. 33pp.
- Thornthwaite, C.W., J.R. Mather, and D.B. Carter. 1958. Three water balance maps of eastern North America. Resources for the Future, Inc. Washington, D.C.
- Thorson, R.M. 2002. Stone by stone: the magnificent history in New England's stone walls. Walker and Company, New York, NY. 287 pp.
- Trani, M.K., R. T. Brooks, T. L. Schmidt, V. A. Rudis, and C.M. Gabbard. 2001. Patterns and trends of early successional forests in the eastern United States. *Wildl. Soc. Bull.* 29(2).
- Trimble, G.R. Jr., J.H. Patric, J.D. Gill, G.H. Moelleer, and J.N. Kochenderfer. 1974. Some options for managing forest land in the central Appalachians. USDA Forest Service, General Technical Report NE-12, 42 pp.
- Trombulak, S.C., and C. Frissell. 2000. A review of the ecological effects of roads on terrestrial and aquatic ecosystems. *Conservation Biology* 14: 18-30.
- Tubbs, C.H., R.M. DeGraaf, M. Yamasaki and W.M. Healy. 1987. Guide to wildlife tree management in New England northern hardwoods. USDA For. Serv. Gen. Tech. Rep. NE-118.
- Tyrrell, L.E. and T.R. Crow. 1994. Structural characteristics of old-growth hemlock-hardwood forests in relation to age. *Ecology* 75(2):370-386.
- Tzipori, S. 1983. Cryptosporidiosis in animals and humans. *Microbiol. Rev.* 47:84.
- U.S. E.P.A. 1993. Handbook: urban runoff pollution prevention and control planning. U.S. publication no. EPA/625/R-93/004.
- U.S.D.A. Forest Service, 2005. B. Onken and R. Reardon, Compilers. Third symposium on hemlock woolly adelgid in the eastern United States. Asheville, North Carolina. February 1-3, 2005. Online: http://www.na.fs.fed.us/fhp/hwa/pub/2005_proceedings/index.shtml
- U.S.D.A. Forest Service. 1991. Riparian forest buffers; function and design for protection and enhancement of water resources. NA-PR-07-91.
- U.S.D.A. Forest Service. 1990. Silvics of North America. Vol.1 Conifers, Vol.2 Hardwoods. Agricultural Handbook 654. Vol.1 675 pp; Vol.2 877 pp.
- United States Geological Survey (USGS). 2000. <http://www.ma.water.usgs.gov/>
- Van Miegroet, H. and D.W. Johnson. 1993. Nitrate dynamics in forest soils. IN: Nitrate: processes, patterns and management. T.P. Burt, A.L. Heathwaite, S.T. Trudgill, eds. John Wiley and Sons, New York, N.Y. 444p.
- Vecellio, G.M., R.D. Deblinger, and J.E. Cardoza. 1993. Status and management of moose in Massachusetts. *Alces*. 29: 1-7.
- Vellidis, G. 1994. Restoration of riparian wetlands: a case study. In: Proceedings: Riparian Buffer Zones Conference. Society of Soil Scientists of Southern New England.
- Vellidis, G., R. Lowrance, and M.C. Smith. 1994. A quantitative approach for measuring N and P concentration changes in surface runoff of a restored riparian wetland forest. *Wetlands* 14(2):73-81.
- Veneman, P.L.M. 1984. Evaluation of the buffering capacity of Massachusetts soils to negotiate the impact of acid deposition on water quality. Completion Report to the Massachusetts Dept. of Environmental Quality Engineering Division of Air Quality control. University of Massachusetts at Amherst.

- Vernegaard, L., R. Hopping, and D. Reid. 1998. Ecological management of grasslands: guidelines for managers. The Trustees of Reservations, Massachusetts. 20pp.
- Verry, E.S. 1986. Forest harvesting and water: the Lakes States experience. *Water Resources Bulletin*. 22(6):1039-1047.
- Vitousek, P. M., W.A. Reiners, J.M. Melillo, C.C. Grier, and J.R. Gosz. 1981. IN: Stress effects on natural ecosystems. pp. 115-127.
- Vitousek, P.M. 1977. The regulation of element concentrations in mountain streams in the northeastern U.S. *Ecol. Monographs*, Vol. 47, pp 65-87.
- Vitousek, P.M. 1985. Community turnover and ecosystem nutrient dynamics. In: Pickett, S.T.A. and P.S. White., eds, *The Ecology of Natural Disturbance and Patch Dynamics*. Academic Press Inc., New York, pp 325-333.
- Vitousek, P.M. and J.M. Melillo. 1979. Nitrate losses from disturbed forests: patterns and mechanisms. *Forest Science* Vol. 25(4).
- Vitousek, P.M. and W.A. Reiners. 1975. Ecosystem succession and nutrient retention: a hypothesis. *Bioscience* 25:376-381.
- Vitousek, P.M., J.R. Gosz, C.C. Grier, J.M. Melillo, and W.A. Reiners. 1982. A comparative analysis of potential nitrification and nitrate mobility in forest ecosystems. *Ecol Monographs* Vol 52(2), pp 155-177.
- Vitousek, P.M., J.R. Gosz, C.C. Grier, J.M. Melillo, W.A. Reiners and R.L. Todd. 1979. Nitrate losses from disturbed ecosystems. *Science*. 204:469-474.
- von Oettingen, S.L. 1982. A survey of beaver in central Massachusetts for *Giardia lamblia*. M.S. Thesis, University of Massachusetts, Amherst. 57 pp.
- Walters, C.J. 1986. Adaptive management of renewable resources. MacMillan Publishing Company, New York, NY.
- Wandle, S.W. n.d. Estimating peak discharges of small, rural streams in Massachusetts. U.S. Geological Survey Water-Supply Paper 2214. 25 pp.
- Waring, R.H. and W.H. Schlesinger. 1985. *Forest ecosystems: concepts and management*. Academic Press, Inc. Harcourt Brace Jovanovich. New York, N.Y.
- Webster, B. 1983. The shy beaver: powerful effects on ecology seen in new research. *N.Y. Times*, Tues. Jan. 11.
- Welsch, D.J. 1991. Riparian forest buffers; function and design for protection and enhancement of water resources. U.S.D.A. Forest Service, Radnot, PA. Pub No. NA-PR-07-91.
- Whigham, D.F., C. Chitterling, and B. Palmer. 1988. Impacts of freshwater wetlands on water quality: a landscape perspective. *Environmental Management*. 12(5):663-671.
- White, L.A., Jr., A.R. Eschner, S.J. Riha, and F.N. Swader. Forest soil and water relationships; New York State forest resources assessment report no. 11. New York State Department of Environmental Conservation, Division of Lands and Forests, Forest Resources Planning Program, 1980.
- Whitney, G.G. 1994. From coastal wilderness to fruited plain: a history of environmental change in temperate North America, 1500 to the present. Cambridge University Press. Cambridge, England. 451 p.
- Wilde, S.A., C.T. Youngberg and J.H. Hovind. 1950. Changes in composition of ground water, soil fertility and forest growth produced by the construction and removal of beaver dams. *J. Wildl. Manage.* 14(2):123-128.

- Wilén, B.O., W.P. McConnell and D.L. Mader. no date. The effects of beaver activity on water quantity and quality. Unpubl. manuscript. Univ. of Mass., Amherst, MA.
- Williams, T.M. and A.C. Mace Jr. 1974. Effects of alternative harvesting systems on cycling of major nutrients in a forest soil. In: Proceedings of the Society of American Foresters.
- Williamson, S. J. n.d.. Forester's guide to wildlife habitat improvement. Univ. New Hampshire, Coop. Ext. 41pp.
- Wischmeier, W.H. and D.D. Smith. 1965. Predicting rainfall-erosion losses from cropland east of the Rocky Mountains. Agr. Handbook 287. USDA, Washington, DC.
- Woo, M-K. and J.M. Waddington. 1990. Effects of beaver dams on subarctic wetland hydrology. Arctic. 43(3):223-230.
- Wraight, S.P., M.A. Jackson, and S.L. deKock. 2001. Production, stabilization and formulation of fungal biocontrols agents, pp. 253-288. In Butt, T.M., C.W. Jackson, and N. Magan (eds.), *Fungi as Biocontrol Agents: Progress, Problems and Potential*. Pp. 253-288. CABI Publishing, New York.
- Wright, R.F. 1976. The impact of forest fire on the nutrient influxes to small lakes in northeastern Minnesota. Ecology. 57:649-663.
- Yeakley, J.A., D.C. Coleman, B.L. Haines, B.D. Kloeppel, J.L. Meyer, W.T. Swank, B.W. Argo, J.M. Deal, and S.F. Taylor. 2003. Hillslope nutrient dynamics following upland riparian vegetation disturbance. Ecosystems (2003) 6:154-167.
- Yeo, J. 1991. The future role of the Sudbury Reservoirs and watershed in the MDC/MWRA water supply system: a study of alternatives. MWRA unpublished report, 54 p.
- Yorks, T.E. 2002. Influence of hemlock mortality on soil water chemistry and ground flora. In Proceedings: hemlock woolly adelgid in the eastern United States. February 5-7, 2002. B. Onken (ed). Rutgers – the State University of New Jersey, East Brunswick, NJ. 403 p.
- Yorks, T.E., J.C. Jenkins, D.J. Leopold, D.J. Raynal, and D.A. Orwig. 2000. Influences of Eastern hemlock mortality on nutrient chemistry. IN:McManus, K.A., K.S. Shields, D.R. Souto, eds. 2000. Proceedings: symposium on sustainable management of hemlock ecosystems in eastern North America; 1999 June 22-24; Durham, N.H. Gen. Tech. Rep. 267. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 237 p.
- Yoshida, S. and M. Ichikuni. 1989. Role of forest canopies in the collection and neutralization of airborne acid substances. The Science of the Total Environment. 84:35-43.
- Yuretich, R.F., ed. 1992. Impacts of acid deposition on watersheds of the Quabbin Reservoir. Water Resources Research Center, University of Massachusetts, Pub. No. 166. 229 pp.
- Yuretich, R.F., P.J. Stekl and S.J. Pohanka. 1986. Hydrogeology and mineral weathering reactions in watersheds of Central Massachusetts. Water Resources Research Center, University of Massachusetts, Pub. No. 152. 85 pp.
- Zhou, G., S. Lui, Z. Li, D. Zhang, X. Tang, C. Zhou, J. Yan, and J. Mo. 2006. Old-growth forests can accumulate carbon in soils. Science. 324:1417

9 Glossary of Terms

Listed in alphabetical order below are terms and definitions that DWSP uses throughout various watershed land management plans. Specific sources of definitions are shown in parenthesis, where applicable.

age class: (from Society of American Foresters, 1971. Terminology of forest science, technology, practice, and products.) one of the intervals, commonly 10 years, into which the age range of tree crops (and sometimes other vegetation) is divided for classification or use.

aggradation: (from Bormann and Likens, 1979. Pattern and process in a forested ecosystem.) in Northern Hardwoods, a period of more than a century when the ecosystem accumulates total biomass reaching a peak at the end of the phase; preceded by the reorganization phase and followed by the transition phase.

advance regeneration: in silvicultural terms, young trees that have become established naturally in a forest, in advance of regeneration cutting; may become established following “preparatory” cuts.

allogenesis: changes in an ecological community primarily through periodic, acute, external (exogenous) disturbances, such as storms. These changes generally reset the successional progression of the community.

area inch; acre inch: used to describe changes in water yield from a given area of land; for instance, if a change in vegetation results in an increase of one acre inch in water yield, this translates to $43,560 \text{ sq ft} \times 1/12 \text{ ft} = 3,630 \text{ cubic feet per acre}$; $3,630 \text{ ft}^3 / 7.5 \text{ gals per ft}^3 = 484 \text{ gallons additional yield per acre}$. Area inch is translated to percent water yield increase by dividing area inch by total inches of yield. For example 40 inches of precipitation generally yields 50%, or 20 inches of discharge, therefore a 2 area inch increase in yield on this watershed is a 10% increase (2/20).

autogenesis: changes in an ecological community primarily through the regular, internal processes of growth, competition, and senescence, which are general endogenous (within community) forces that result in a steady successional progression of the community.

basin; sub-basin: the land area from which all water flows to a single, identified water source, such as a stream, a river, or a reservoir. Sub-basin is used to refer to the basin of a tributary or lower *order* stream (the higher the order, the greater the area drained).

basal area: the area in square feet of the cross section of a tree taken at 4.5 feet above the ground.

“beaver pipe”; flow control pipe: generally a length of culvert that is extended into a beaver pond and at or near the top of the beaver dam, in order to maintain the pond level at a particular level.

Best Management Practices, BMPs: in natural resources management, refers to a set of standards that have been designed for an activity, and often a region, to protect against degradation of resources during management operations.

biological diversity (biodiversity): a measure, often difficult to quantify, of the variety and abundance of plant and animal species within a specified area, at the genetic, species, and landscape level of analysis. The 1992 UN Convention on Biological Diversity defined biodiversity as “the variability among living organisms from all sources including, inter alia, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystems”.

biomass: (SAF) the total quantity of living organisms of one or more species per unit area (species biomass) or of all the species in a community (community biomass)

Conservation Management Practices, CMPs: in natural resources management, refers to a set of standards that have been designed for an activity, and often a region, to protect against degradation of resources during management operations. Also known as Best Management Practices, BMPs.

conservation restriction; conservation easement; CR: a legal agreement between a landowner and another party whereby the landowner deeds the rights to development of the property to the other party, but retains ownership of the land and other rights to its use. Specific agreement varies, but the general result is to protect land from conversion to new uses without requiring transfer of ownership; DWSP also limits or retains the right to approve certain agricultural and silvicultural practices in its CRs.

Continuous Forest Inventory (C.F.I.): an extensive method of forest inventory in which permanent sample plots are remeasured at periodic intervals to determine forest growth and condition; DWSP's CFI is composed of 1/5 acre permanent plots, located on a 1/2 mile grid, and remeasured every 10 years.

cutting cycle: the frequency with which silvicultural cuttings are conducted in any given area; cutting cycle is a subunit of "rotation," which is determined either by the maximum life of the existing overstory, or by a predetermined maximum age imposed on the area.

Cryptosporidium: A coccidian protozoan parasite found in humans and various wild and domestic animals that can be transmitted via water and often causes serious intestinal illness. While the epidemiology and transmission of *Cryptosporidium* are similar to *Giardia*, its oocysts are smaller than the cysts of other protozoa, and thus may be more difficult to remove from water supplies.

diameter at breast height; DBH: the diameter of a tree, outside the bark, taken at 4.5' above the ground, generally in inches and fractions.

diverse/diversity: in this plan, the term is most often used to refer to forest composition, and refers to both height or size diversity in trees, seeking a minimum of three distinct layers (understory, midstory, and overstory), and to diversity of species composition, with a general goal of avoiding monocultures and working to include components of hemlock, pine, oak, birch, and maple throughout the forest.

disturbance-sheltered: areas that based on slope and aspect are physically "sheltered" from the influence of a catastrophic New England hurricane blowing from the southeast, based on a model developed at the Harvard Forest; the most sheltered areas are steep slopes facing northwest.

edge effect: this term has traditionally been used to describe the increased richness of flora and fauna found where two habitat types or communities meet. More recently, the term has also been used to refer to the increased predation and brood parasitism that often occurs near these boundaries.

endogenous disturbance: disturbance that originates within the ecological community. For example, a single tree that succumbs to a root-rot fungus and falls to the ground, breaking off several other trees on the way, creates an endogenous disturbance. While the proximal cause of the treefall may be wind or accumulation of snow and ice, the primary cause is still considered endogenous in this instance. (see exogenous disturbance below)

even-aged: (SAF) an area of forest composed of trees having no, or relatively small, differences in age. NOTE: By convention the maximum difference admissible is generally 10 to 20 years, though with rotations of 100 years or more, differences up to 30% of the rotation may be admissible.

exogenous disturbance: disturbance that originates from forces outside of the ecological community. For example, storms that carry high winds can cause large-scale treefall well in advance of normal senescence and decay. The cause of the disturbance is therefore considered exogenous. (see endogenous disturbance above)

feller-buncher; feller-buncher-processor: logging machines that grasp a tree to be cut or “felled,” sever it at the stump with either a saw or hydraulic shears, and directionally drop it to the ground. Some machines can accumulate, or “bunch” several trees before releasing them. The most complex machines are also capable of delimiting and sawing trees into predetermined lengths (processing).

forest canopy: (SAF) the more or less continuous cover of branches and foliage formed collectively by the crowns of adjacent trees and other woody growth.

forest fragmentation: the separation of a previously contiguous forested area into discontinuous patches or “fragments.” These fragments are less useful to wildlife that require large contiguous habitats. Fragmentation by suburban development is likely to be detrimental to “deep woods” species, while the simple break imposed by an access road is not often an impediment.

forwarder: a logging machine used to “forward” logs from the woods to a landing. A forwarder differs from a skidder in that the logs are hydraulically loaded onto the machine and carried, rather than skidded through the woods.

G.I.S.: Geographic Information System - a computer-based analysis and mapping system for spatially-linked data sets.

Giardia: A protozoan parasite found in humans and various wild and domestic animals that can be transmitted via water and often causes serious intestinal illness.

group selection, or small group selection: a cutting in which a roughly homogenous group of overstory trees is removed at the same time, and regeneration is either initially established or released as a result. Groups that, when cut, leave openings with diameters that exceed twice the height of the surrounding trees (and therefore contain areas not under the influence of those trees) are generally referred to as “patches”.

hurricane exposure (“exposed,” “intermediate,” “sheltered”): generally used in DWSP management plans to mean physical exposure of a site to catastrophic hurricane winds, those coming from the southeast. Research at the Harvard Forest in Petersham, MA provides a model of the impact of this typical New England hurricane, which includes slope and aspect. Actual damage will depend on the type and size of vegetation present.

intermediate cut: cutting of trees in a stand during the period between establishment and maturity. Objectives may include the improvement of vigor by reducing competition or the manipulation of species composition. Regeneration may occur following intermediate cuts, but it is incidental to the objective of intermediate cutting.

irregular shelterwood: similar to the shelterwood silvicultural system except that overstory removals are protracted, taking as long as half the rotation, so that the resulting new stand is quite uneven-aged (wide intervals between the oldest and youngest trees) and mimics the multi-storied effect of strictly uneven-aged systems.

log landing; landing: a clearing of variable size to which logs, pulp, and/or firewood are skidded or forwarded during a logging operation, in order to facilitate their processing or further transport by truck.

mast: the fruit and seeds of trees and shrubs. Mast constitutes an important food source for many wildlife species.

milacre: one-thousandth of an acre.

mineral soil: any soil consisting primarily of minerals (sand, silt, and clay), rather than organic matter.

multi-aged: a forest that contains more than a single age class and may contain many different age classes not necessarily in balanced proportions. The absence of strict balancing of age classes distinguishes these forests from true “uneven-aged” forests.

multi-storied or multi-layered forest: a forest containing a distinct understory, midstory, and overstory. From a watershed perspective, these layers provide, respectively, immediate response to disturbance, vigorous uptake of nutrients, and deep filtration of air-borne and precipitative pollutants.

naturally managed: the results of a deliberate decision to allow natural disturbances and processes prevail by adopting a minimal management approach that protects forests from development or other land use changes and possibly human-caused fire, but which includes vegetation management only where it clearly counteracts a negative result from previous human disturbances.

old-growth: various definitions, but one definition in Massachusetts is that old-growth must contain at least four acres in which the dominant trees are late-successional species, have reached at least half the potential maximum age of the species with a few trees at or near that age (e.g., birches/maples at 300 years or hemlocks at 400), have late-successional tree regeneration present, and show little or no evidence of human or large-scale natural disturbance during the current stand’s development.

patch cut: a regeneration opening in the forest of various sizes but in which some part of the opening is not under the influence of the surrounding trees. All of the trees in the designated area are cut. A patch cut is a small clearcut, but one which removes only a fraction of a complete stand, resulting in a multi-aged stand once the opening has regenerated.

preparatory cutting: (SAF) removing trees near the end of a rotation so as to open the canopy and enlarge the crowns of seed bearers, with a view to improving conditions for seed production and the establishment of natural regeneration.

protected: refers to areas of the watershed that, according to the Harvard Forest model of hurricane disturbance, would suffer minimal damage from the recurrence of a hurricane similar to that of 1938, due primarily to topography and orientation.

protection forest: (SAF) an area, wholly or partly covered with woody growth, managed primarily to regulate stream flow, maintain water quality, minimize erosion, stabilize drifting sand, or to exert any other beneficial forest influences

regeneration: recently established tree growth, generally smaller than one inch dbh; also, the process of establishing this growth, as in “bring about the regeneration of a forest area”.

regeneration cut: (SAF) any removal of trees intended to assist regeneration already present or to make regeneration possible.

riparian: pertaining to the bank of a stream or other water body; (SAF) vegetation growing in close proximity to a watercourse, lake, swamp, or spring, and often dependent on its roots reaching the water table.

rotation: in conventional forestry, rotation is (SAF) the planned number of years between the formation or regeneration of a crop or stand and its final cutting at a specified stage of maturity. In the selection system of uneven-aged management, however, the concept of a rotation is replaced with the average age of trees removed to initiate regeneration.

salvage; salvage cutting: the removal of trees damaged by fire, wind, insects, disease, fungi, or other injurious agents before their timber becomes worthless. In some situations, the motivation is the reduction of fuel loading and fire hazard. Sanitation cutting is related, but is a proactive removal of diseased or highly susceptible trees in order to slow or halt the spread of a disease or other destructive agent.

seep: a wet area, generally associated with groundwater seepage, which is important to wildlife because it remains unfrozen, and generally uncovered, during periods when the ground is otherwise snow-covered, which makes it easier for wildlife to forage for seeds.

shelterwood: (SAF) mostly even-aged silvicultural systems in which, in order to provide a source of seed, protection for regeneration, or a specific light regime, the overstory (the shelterwood) is removed in two or more successive shelterwood cuttings, the first of which is ordinarily the seed cutting (though it may be preceded by a preparatory cutting) and the last of which is the final cutting, while any intervening cuttings are termed removal cuttings. Note that where adequate regeneration is already present, the overstory may be removed in one cutting, resulting in a method referred to as a one-cut shelterwood.

silviculture: (SAF) generally, the science and art of cultivating (i.e., growing and tending) forest crops, based on a knowledge of silvics (the study of the life history and general characteristics of forest trees and stands, with particular reference to environmental factors affecting growth and change). More particularly, silviculture is the theory and practice of controlling the establishment, composition, constitution, and growth of forests.

site: in forestry, the combination of environmental factors that affect the ability of a species to grow and persist, including at least soil characteristics, aspect, altitude and latitude, and local climate. Sites are often classified by the ability of specific trees to grow on them.

site index: the ability of a given site to grow a given species. As height growth is generally not density dependent, a common forestry site index is the height to which a given species will grow on the site in fifty years (so that a site with a Red Oak site index of 65 will grow Red Oak to that height in fifty years).

site preparation: in silviculture, any of a variety of treatments of a site that are intended to enhance regeneration success. A common goal of these treatments is to remove enough of the accumulated organic layers above the mineral soil so as to expose that soil and enhance the ability of seeds that fall on it to germinate and grow. The simple skidding of logs is an incidental, and often sufficient, site preparation.

site-suited: species that have evolved to take advantage of a particular type of site. Where species are planted on other sites, they may succumb prematurely to disturbance or disease. Red pine grows and persists well on deep, sandy soils, where root rots are less common, but may become excessively prone to wind and or root rotting diseases on the moist agricultural soils on which they were typically planted.

skidder: logging machine used to “skid” logs from the woods to a landing or a forwarder road. Logs are either winched by cable to the skidder (cable skidder), or lifted on one end by a hydraulic grapple (grapple skidder), and then dragged.

stand: (SAF) a community of trees possessing sufficient uniformity as regards composition, constitution, age, spatial arrangement, or condition to be distinguishable from adjacent communities.

steady state: (Bormann and Likens, 1979. p.4) “For the ecosystem as a whole, over a reasonable period of time gross primary production equals total ecosystem respiration, and there is no net change in total standing crop of living and dead biomass”.

stocking: in forestry, the extent to which a site is occupied by trees compared to the maximum occupation possible at a given stand age; a relative measure of stand density. Most commonly measured as basal area per acre, stocking is often related directly to crown closure, as a site is considered fully occupied when crown closure is complete (when each crown has grown to touch all adjacent ones). As crowns can be of very different sizes among species and tree ages within stands, average diameter (dbh) and total number of trees of a “fully stocked” site is variable.

stream order: a classification of streams within watersheds. Small streams at the uppermost level of stream systems are labeled “first-order”; two first-order streams join to form a “second-order” stream; two second-order streams join to form a “third-order” stream; etc.

succession: (SAF) the gradual supplanting of one community of plants by another, the sequence of communities being termed a “sere” and each stage “seral.” Succession is “primary” (by “pioneer species”) on sites that have not previously borne vegetation, “secondary” after the whole or part of the original vegetation has been supplanted; “allogenic” when the causes of succession are external to and independent of the community (e.g., a storm, or climate change), and “autogenic” when the developing vegetation is itself the cause. “Early succession” generally refers to the pioneer stages and species that follow disturbance, while “late succession” refers to stages and species that occur as an area continues to develop undisturbed for long periods.

thinning: an intermediate silvicultural treatment, generally with the goal of altering the forest composition and/or improving the growing conditions for the residual trees, regardless of associated regeneration effects. Most thinnings leave stands considered to be fully stocked, i.e., capable of fully occupying the site a short while after the thinning has been completed.

turbidity: a water quality measure that is most commonly derived by measuring the proportion of a given amount of light that is deflected by suspended/dissolved sediments in a water sample, giving an indirect measure of these sediments. Most common unit is the nephelometric turbidity unit, NTU.

uneven-aged: (SAF) a forest, crop, or stand composed of intermingling trees that differ markedly in age. By convention, a minimum difference between tree ages of 25% of the rotation age is generally accepted. Some texts require a minimum of three distinct age classes for a stand to qualify as “uneven-aged.” Uneven-aged silviculture, when fully applied, results in balanced age classes. Mixed-age class forests in which no attempt is made to balance age classes is referred to simply as “multi-aged forest”.

vernal pool: a temporary body of fresh water that provides crucial habitat for several vertebrate and many invertebrate species of wildlife, but does not support fish populations.

wetland: generally refers in DWSP land management plans to areas defined as “wetlands” by M.G.L. C.131 § 40 (the “Wetlands Protection Act”) and 310 C.M.R. 10.00 (the “Wetlands Protection Regulations”), updated as these are revised.