

Lyme Disease and Babesiosis Co-Infection*:2009-2013

Massachusetts Department of Public Health

*Co-infection = patient with confirmed/probable Lyme disease + laboratory evidence of *B. microti* infection.
Symptomatic co-infection = patient with confirmed/probable case status for both Lyme disease and babesiosis


2009-2013 Surveillance Highlights

- Between 2009 and 2013, there were 24,687 confirmed and probable cases of Lyme disease, and 1,084 confirmed and probable cases of babesiosis reported in Massachusetts.
- There were 294 *Borrelia burgdorferi/Babesia microti* co-infections identified in Massachusetts; 171 of these patients met the case definition for both diseases (symptomatic co-infection).
- The highest incidence rates of *B. burgdorferi/B. microti* co-infection occurred in Dukes and Nantucket counties (67 and 197 per 100,000 residents). Some risk of co-infection exists throughout Massachusetts.
- The majority of *B. burgdorferi/B. microti* co-infections occurred in June, July and August.
- Patients with Lyme disease and babesiosis symptomatic co-infection were more likely to be reported with joint pain, muscle pain, headache and cardiac manifestations than patients with either Lyme disease or babesiosis alone.
- 36% of patients with Lyme disease and babesiosis symptomatic co-infection were hospitalized, which is similar to 39% of patients with babesiosis alone.
- 35% of reported babesiosis cases in children less than 10 years old had Lyme disease symptomatic co-infection.
- 45% of *B. burgdorferi/B. microti* co-infections occurred in 50 - 69 year olds.
- 15% of patients 50 years and older with confirmed or probable babesiosis had Lyme disease symptomatic co-infection; 1% of patients in this age group with confirmed or probable Lyme disease had babesiosis symptomatic co-infection.

MAP: Reported babesiosis cases overlaid on reported Lyme disease cases to demonstrate areas with higher risk of co-infection. (NOTE: these cases are reported by city/town of residence, not necessarily place of exposure).


COMPARE TO

TABLE: Actual counts and incidence rates of patients with confirmed or probable Lyme disease and laboratory evidence of *B. microti* infection.



County	<i>B. burgdorferi/B. microti</i> co-infection (N=294)	Co-infection 5 year incidence rate (per 100,000)
Barnstable	65	30
Berkshire	1	1
Bristol	20	4
Dukes	11	67
Essex	34	5
Franklin	1	1
Hampden	12	3
Hampshire	3	2
Middlesex	45	3
Nantucket	20	197
Norfolk	23	3
Plymouth	48	10
Suffolk	5	1
Worcester	6	1

Figure 2: Number of confirmed and probable cases of babesiosis and cases of babesiosis with Lyme disease symptomatic co-infection in Massachusetts by age: 2009-2013



- The greatest proportion of babesiosis cases with Lyme disease symptomatic co-infection occurred in children less than 10 years of age (35%).
- A large number of the symptomatic co-infection cases occurred in the 50 - 79 year age range.
- 15% of patients 50 years and older, with confirmed or probable babesiosis had Lyme disease symptomatic co-infection. Only 1% of patients in this age group, with confirmed or probable Lyme disease had babesiosis symptomatic co-infection.

Table 3: Percent of particular reported signs and symptoms among probable and confirmed cases of Lyme disease, babesiosis, and Lyme disease/babesiosis symptomatic co-infection reported in Massachusetts: 2009-2013

Sign or Symptom	Lyme (n=24,687) Percent	Babesiosis (n=1,084) Percent	Lyme/Babesiosis Symptomatic co-infection (n=171) Percent
Rheumatologic	55	47	75
Arthritis	23	0	12
Joint pain	43	46	58
Neck pain	14	2	12
Cardiac	2	0	5
Fatigue	36	79	82
Fever	34	83	85
Gastrointestinal	1	44	37
Muscle Pain	42	50	58
Headache	25	50	59
Neurological	14	1	14
Cognitive impairment	5	0	8
Erythema migrans	49	1	39

- Reported signs and symptoms of Lyme disease include arthritis; cardiac abnormalities such as arrhythmia, AV block and palpitations; neurological (auditory impairment, Bell’s palsy, lymphocytic meningitis, paresthesia, radiculoneuropathy, and visual and cognitive impairment); rash (including erythema migrans); and headache.
- Reported signs and symptoms of babesiosis include joint pain and muscle ache; fatigue and malaise; fever (fever, chills and sweats); headache; and gastrointestinal symptoms (anorexia and nausea).
- **Reported cases with Lyme disease and babesiosis symptomatic co-infection were more likely to experience joint pain, muscle pain, headache and cardiac manifestations than patients with either Lyme disease or babesiosis alone (p<0.05 for each of these signs or symptoms).**